

Valerie K. Brown

BIOLOGICAL JOURNAL

of the Linnean Society

CABI BIOSCIENCE
EGHAM LIBRARY

VOLUME 55

NUMBERS 1-4

PAGES 1-386

THE LINNEAN SOCIETY OF LONDON
BURLINGTON HOUSE, PICCADILLY, LONDON W1V 0LQ
1995

Dates of publication of the numbers included in Vol. 55:

No. 1, pp. 1–92, 31 May 1995

No. 2, pp. 93–182, 20 June 1995

No. 3, pp. 183–272, 10 July 1995

No. 4, pp. 273–386, 21 August 1995

CONTENTS

	Pages
BARNARD, P., Timing of ornament growth, phenotypic variation, and size dimorphism in two promiscuous African whydahs (Ploceidae: <i>Vidua</i>)	129–141
CHEVILLON, C., ADDIS, G., RAYMOND, M. and MARCHI, A., Population structure in Mediterranean islands and risk of genetic invasion in <i>Culex pipiens</i> L. (Diptera: Culicidae)	329–343
EMBERTON, K. C., Cryptic, genetically extremely divergent polytypic, convergent, and polymorphic taxa in Madagascan <i>Tropidophora</i> (Gastropoda: Pomatiastidae)	183–208
FITZPATRICK, S., BERGLUND, A. and ROSENQVIST, G., Ornaments or offspring: costs to reproductive success restrict sexual selection processes	251–260
FORSMAN, A. and SHINE, R., The adaptive significance of colour pattern polymorphism in the Australian scincid lizard <i>Lampropholis delicata</i>	273–291
GASTON, K. J., SCOBLE, M. J. and CROOK, A., Patterns in species description: a case study using the Geometridae (Lepidoptera)	225–237
JORMALAINEN, V., MERILAITA, S. and TUOMI, J., Differential predation on sexes affects colour polymorphism of the isopod <i>Idotea baltica</i> (Pallas)	45–68
KIRSCH, J. A., LAPOINTE, F.-J. and FOESTE, A., Resolution of portions of the kangaroo phylogeny (Marsupialia: Macropodidae) using DNA hybridization	309–328
LYNCH, J. M. and HAYDEN, T. J., Genetic influences on cranial form: variation among ranch and feral American mink <i>Mustela vison</i> (Mammalia: Mustelidae)	293–307
MALLET, J. and GILBERT, L. E. Jr., Why are there so many mimicry rings? Correlations between habitat, behaviour and mimicry in <i>Heliconius</i> butterflies	159–180
MARPLES, N. M. and BRAKEFIELD, P. M., Genetic variation for the rate of recruitment of novel insect prey into the diet of a bird	17–27
MEDEL, R. G., Convergence and historical effects in harvester ant assemblages of Australia, North America, and South America	29–44
NYLIN, S., WICKMAN, P. O. and WIKLUND, C., Life-cycle regulation and life history plasticity in the speckled wood butterfly: are reaction norms predictable?	143–157
PETANIDOU, T., ELLIS, W. N. and ELLIS-ADAM, A. C., Ecogeographical patterns in the incidence of brood parasitism in bees	261–272
PETERSON, M. A., Unpredictability in the facultative association between larvae of <i>Euphilotes enoptes</i> (Lepidoptera: Lycaenidae) and ants	209–224
PRENTER, J., MONTGOMERY, W. I., and ELWOOD, R. W., Multivariate morphometrics and sexual dimorphism in the orb-web spider <i>Metellina segmentata</i> (Clerck, 1757) (Araneae, Metidae)	345–354
SHERRATT, T. N. and MACDONALD, A. D., Some population consequences of variation in preference among individual predators	93–107
URIZ, M. J. and MALDONADO, M., A reconsideration of the relationship between polyaxonid and monaxonid spicules in Demospongiae: new data from the genera <i>Crambe</i> and <i>Discorhabdella</i> (Porifera)	1–15
WARD, L. K., HACKSHAW, A. and CLARKE, R. T., Food plant families of British insects and mites: the influence of life form and plant family	109–127
WEINTRAUB, J. D., LAWTON, J. H. and SCOBLE, M. J., Lithinine moths on ferns: a phylogenetic study of insect-plant interactions	239–250
WILSON, P., Selection for pollination success and the mechanical fit of <i>Impatiens</i> flowers around bumblebee bodies	355–383
YU, H. T., Patterns of diversification and genetic population structure of small mammals in Taiwan	68–69
 REVIEWS	
ANDERSSON, M., <i>Sexual Selection</i> (M. Andersson)	92
BENTON, M. J. and SPENCER, P. S., <i>Fossil Reptiles of Great Britain</i> (F. M. Braodhurst).	385–386
BOWLES, M. L. and WHELAN, R. J., <i>Restoration of Endangered Species, Conceptual Issues, Planning and Implementation</i> (N. W. Moore).	181

GILBERT, J., DANIELPOOL, D. L. and STANFORD, J. A., <i>Groundwater Ecology</i> (A. A. Savage) .	385
LEVER, C. and POYSER, A. D., <i>Naturalised Animals: the Ecology of Successfully Introduced Species</i> (D. W. Yalden)	91
MADDOCKS, L., BONE, Q and RAYNOR, J. M. V., <i>Mechanics and Physiology of Animal Swimming</i> (H. D. Jones)	385
SOLBRIG, O. T., VAN EMDEN, H. M. and VAN OORDT, P. G. W. J., <i>Biodiversity and Global Change</i> (O. L. Gilbert)	181-182
STEVENS, P. F., <i>The Development of Biological Systematics</i> (D. Williams)	386

INDEX

compiled by Stanley Thorley

Abbreviations:

App. – Appendix

Fig. – Figure

n. – note

Tab. – Table

*signifies a keyword.

- A2–B2 (resistance genes): *Culex pipiens* population structure and genetic invasion 329–331, 340–341
- AAT (aspartate aminotransferase)
- Culex pipiens* population structure and genetic invasion 331
- Aat* locus 332 Tab. 1, 337–338 Tabs 3–4
- Madagascan *Tropidophora* (allozyme data)
- Aat-1* 186, 189 Fig. 2, 205–206 App.
- Aat-2* 186, 189 Fig. 2, 205–206 App.
- Abies* spp.: Reecer Creek 211–212
- Academy of Natural Sciences of Philadelphia (ANSP):
- Madagascan *Tropidophora* 201, 205–208 App
- Acanthaceae (*Blechum*): host to *Siproeta stelenis* (nymphalid butterflies) 240
- acanthostyles: spicules in Demospongiae 6, 11–12
- asterose 1, 3–4 & Fig. 1, 8 Fig. 5, 11–12
- tuberosa 1, 5–6
- Acarnus*: spicules in Demospongiae 2
- acavid snails: Madagascar 186
- Aceraceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 125 App. 1, 127 App. 2
- ACO (aconitate hydratase): *Culex pipiens* population structure and genetic invasion 331
- Aco* locus 332 Tab. 1, 337–338 Tabs 3–4
- Aco-Fum* pair 336
- Aconitase (*Acon-1*, *Acon-2*, E.C. 4.2.1.3) 72
- Acon-1*
- Apodemus* 87 App. 2
- Microtus* 89 App. 2
- Niviventer* 87 App. 2
- Acromyrmex*: South American harvester ants 34, 40
- A. aspersus* 44 App.
- A. lobicornis* 44 App.
- actines: spicules in Demospongiae 1–3, 5–6 & Fig. 3, 8, 11
- see also monactines
- Adalia bipunctata* (two-spot ladybirds) 17–19
- black form with four or six red spots (melanic) 19–20 & Fig. 1, 24 Tab. 2, 26
- red form with two black spots 17–20 & Fig. 1, 22–23 & Tab. 1, 26
- adaptation: *Pararge aegeria* (speckled wood butterflies) 143–144, 153
- adiantoid ferns: dennstaedtioid lineage 248
- Aepyrymus*: resolution of kangaroo phylogeny using DNA hybridization 309, 316–317 Tab. 2, 319–321 & Fig. 2, 323
- A. rufescens* (rufous rat-kangaroo) 309, 313–317 & Tabs 1–2 & Fig. 1, 320–321 Fig. 2
- *aerodynamics: whydahs (*Vidua*) 129, 136, 138–139
- Aethia cristatella* (crested auklets): sexual selection acting on females 256 Tab. 1, 258
- Africa
- Afrotropics: biogeographic region (4) 226
- North: Western Palaearctic (3) 226
- whydahs (*Vidua*) 129–130
- see also Namibia
- Afrotropics
- biogeographic region (4) 226, 228–231 & Tabs 1–2 & Fig. 3, 234 Fig. 8
- patterns in brood parasitism 271 App.
- *age dimorphism: whydahs (*Vidua*) 129–133 & Tab. 1
- *aggregation: unpredictable variation in a facultative mutualism 209–216
- Agraulis vanillae* (butterflies): 'orange' mimicry rings 165
- Alauda arvensis* (skylark) males: flight display compensations 138
- Albany County, New York: fit of flowers around the bodies of bees 361 & Tab. 1, 368, 370 & Fig. 7, 373–376 Tabs 5–8
- Alboran Island, Spain: Demospongiae 3, 7 Fig. 10
- albumin (*Alb*, E.C. 4.2.1.3) 72
- Niviventer* 88 App. 2
- algae
- green (*Cladophora* sp.): juvenile isopod *Idotea baltica* 45, 48–50, 52–55 & Fig. 2 & Tab. 2, 62–63
- insect food-plants 111, 113–118 & Figs 1–3, 125 App. 1, 127 App. 2
- Alismataceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Alimatales 120
- Alismatidae 113
- alleles
- Asian small mammals 69, 72–75 & Tabs 1–2, 82, 87–89 App. 2
- 'private alleles' 73, 77 & Tab 4, 82
- coding secondary traits 139
- colour patterns in a lizard 274
- Culex pipiens* population structure and genetic invasion 331–333 & Tab. 1, 335–338 & Tab. 3
- isopod *Idotea baltica* phenotypes 49
- Madagascan *Tropidophora* 183, 186, 189 & Fig. 2, 200, 205–208 App.
- *allocation theory: sexual selection acting on females 251–253, 255–258 & Fig. 2 & Tab. 1
- *allozyme cladistics: Madagascan *Tropidophora* 183–184, 186–190 & Tab. 1 & Fig. 2, 200
- see also data
- *allozyme variation: Asian mammals 69, 71, 73–75, 78, 80 & Fig. 3, 82
- allozymes
- Culex pipiens* population structure and genetic invasion 331

evolutionary genetics of Asian mammals 69, 80 &

Fig. 3

Impatiens capensis survey 358

Madagascan *Tropidophora* 183, 189 Fig. 2, 200, 205–208 App.

see also *Aat-1*, *Aat-2*, *Est-1*, *Est-2*, *Gbp*, *Gpi*, *Lap*, *Mdh*, *Mpi*, *Pep-G*, *Pep-T1*, *Pep-T2*, *6Pgd*, *Pgm*, *Sod*, *Sordh*, *Tc-6* and *Teb-9*

Alnus (Betulaceae): *Tacpara* associations 247, 250 App.

alpha diversity: convergence in harvester ant assemblages 34, 36 & Tab. 2

Alphitobius diaperinus (dark brown lesser mealworms): insect prey experiments 19

Alsophila (Cyatheaceae: tree ferns): moth associations 242, 244–245 Tab. 2 & Fig. 2, 250 App.

Amaryllidaceae: insect food-plants 114 Fig. 1, 120–121, 125 App. 1, 127 App. 2

America see Central, North and South

Amphipoda (*Gammarus pulex*): *Notonecta* prey 104–105 & Tab. 1

Amphipyrrinae (lepidopteran subfamily): fern associations 241 Tab. 1

analyses

behaviour and mimicry in *Heliconius* 163

colour patterns in a lizard 276–278, 281–283

see also ANCOVA, CATMOD, log-linear, regressions, MANOVA and PHYLIP

CONSENSE 312

convergence in harvester ant assemblages 29–31, 33–39 & Tabs 2–3 & Figs 2–4

pairwise-continent 35–36 & Tab. 2, 38–39

three-continent 35–38 & Figs 2–3 & Tabs 2–3

see also ANOVAs

cranial variation in mink 296, 298–303 & Figs 1–4 CATMOD 333–334

see also ANOVAs, canonical variate (CVA) and multiple group principal components (MGPCA)

Culex pipiens population structure and genetic invasion 335–336 & Tab. 2, 341

LINKDOS 335

Ohta indices 335–336

evolutionary genetics of Asian mammals 70–73, 80–81, 85–86 App. 1

BIOSYS-1 72

PAUP 72–73, 78

phylogenetic 69–71, 78, 80–81

insect food-plants, life form and taxonomy 109–111, 113, 115–117, 118–119, 121–123

see also GENSTAT, multivariate and principle coordinates (PCA)

lithinine moths on ferns: PAUP 246

Madagascan *Tropidophora* 183–184, 186, 200–201 allozyme 183–184

BIOSYS-1 186

cladistic 186

Hennig86 186, 189 Fig. 2

ornament growth, variation and dimorphism in *Vidua* 131–132

StatGraphics 132

patterns in brood parasitism 262–263

see also ANOVAs

resolution of kangaroo phylogeny using DNA

hybridization 309, 311–313, 319–326 & Fig. 2 cladistic 319

PHYLIP 312

sexual dimorphism in *Metellina segmentata* 345–350 discriminant 349–350

see also multiple (MGPCA), multivariate, principle components (PCA) and univariate

sexual selection acting on females 252, 257

unpredictable variation in a facultative mutualism 213

variance for *Impatiens* pollen production 363–364 & Tab. 4, 372–377 & Tabs 6–8

see also meta-analysis and statistical procedures

ANCOVAs: colour pattern polymorphism in a lizard 276–277, 281

Andersson, M. *Sexual Selection*. (review: P. N. Ferns) 92 Andes

Patagonia: Lithinini 241

South America: separation of harvester ant assemblages 39

androecia: fit of flowers around the bodies of bees

357–358, 360–361, 363–367 & Figs 2–4,

369–370, 373 Tab. 5, 375–378 & Tab. 8

see also anthers

Angiospermae

insect food-plants 110, 114–115, 117–121 & Figs 4–8

larval hosts for Lithinini 250 App.

animals

fit of flowers around their bodies 356–357, 378, 380

see also bees, birds, insects and moths

sexual selection acting on females 258–259

lekking 256 Tab. 1

see also birds, pipefish, seahorses and sticklebacks

annuals: insect food-plants 109, 111, 116, 119, 121–123

Anolis sagrei: female colour pattern polymorphism 274 ANOVAs

cranial variation in mink 296–297 & Tab. 2

fit of flowers around the bodies of bees 363–364 & Tab. 4, 368–369, 376 Tab. 8

patterns in brood parasitism 262–263

polymorphism of the isopod *Idotea baltica* 52, 61

*ANOVAs (analyses of variance): convergence in harvester ant assemblages 29–31, 34–35, 39, 41

Schluter's procedure 30, 40–41

ANSP see Academy of Natural Sciences of Philadelphia

Anthelia: citation as *Thallopaga* 250 App.

anthers: fit of flowers around the bodies of bees 355, 357, 360, 366, 372, 378–380

Anthomyiidae: phytophagous insects 110–111

*anti-apostatic selection: variation in predator preference 93–107 & Figs 1–6 & Tab. 1

ants

harvester: convergence and historical events 29–44 & Figs 1–4 & Tabs 1–2 & App.

see also

Australia: *Chelaner*, *Melophorus*, *Meranophus*, *Monomorium*, *Pheidole* and *Tetramorium*

North America: *Messor*, *Novomessor*, *Pheidole*, *Pogomyrmex* and *Solenopsis*

South America: *Acromyrmex*, *Araucomyrmex*, *Brachymyrmex*, *Dorymyrmex*, *Elasmophheidole*, *Forelius*, *Pheidole*, *Pogonomymyrmex*, *Premolepis*, *Pseudomyrmex* and *Solenopsis*

- unpredictable variation in a facultative mutualism
209–223 & Tabs 1–5 & Figs 1–4
ant-larval associations 214–220 & Figs 1–4 &
Tabs 2–5
- Aphalaridae-Trioizidae (psyllids): phytophagous insects
111
- aphids
fireweed and ants 210, 214
phytophagous insects 110–111, 115
calliphorid 115
Mindaridae 111
- Apocynaceae: insect food-plants 112 Tab. 1, 114 Fig.
1, 125 App. 1, 127 App. 2
- Apodemus*: Asian mammals 69–71, 73–76 & Tab. 2,
78–81 & Figs 2–3
A. agrarius 70, 72–73, 76, 78–80 & Figs 2–3, 85
App. 1, 87 App. 2
A. semotus 69–82 & Tabs 1–4 & Figs 2–3, 85 App.
1, 87 App. 2
- *Apoidea: patterns in brood parasitism 261–262,
270–272 App.
see also bees
- aposematism
genetics of predation 17–19
variation in predator preference *see* anti- and pro-
apostatic selections
- *apostatic selection: polymorphism of the isopod
Idotea baltica 45–47, 61–63, 65–66
- Aquifoliaceae: insect food-plants 114–115 & Fig. 1,
125 App. 1, 127 App. 2
- Araceae: insect food-plants 114 Fig. 1, 116, 125 App.
1, 127 App. 2
- Araliaceae: insect food-plants 112 Tab. 1, 114–115 &
Fig. 1, 125 App. 1, 127 App. 2
- Araliales 112
- Araneae: sexual dimorphism in *Metellina segmentata*
345
- Araucomyrmex*: South American harvester ants 40
A. goetschi 44 App.
A. tener 44 App.
- Archicarinae: patterns in species description 226,
232–233 & Tab. 4
- Arctiidae: pteridophyte associations 240
- Argentina: harvester ant assemblages 31–34 & Fig. 1,
39–40, 44 App. & ns
see also Monte desert
- Argyrolagidae (marsupials): Pleistocene 41
- *arid ecosystems: patterns in brood parasitism 261,
267
- arthropods: pteridophagy 240
- Asellus aquaticus* (Crustacea: Isopoda): *Notonecta* prey
104–105 & Tab. 1
- *Asian mammals: evolutionary genetics 69–89 & Figs
1–3 & Tabs 1–4 & Apps 1–2
see also Borneo, China, Hainan, Java, Korea, Suma-
tra and Taiwan
- Aspidium* (*Polystichum*): *Gonophylla* host 250 App.
- Asteridae: insect food-plants 119
- Astrosporidia: polyaxonid and monaxonid spicules in
Demospongiae 12
- auklets, crested (*Aethia cristatella*): sexual selection
acting on females 256 Tab. 1, 258
- Australia
biogeographic region (8) 227–231 & Tabs 1–2 &
Fig. 3, 234–235 & Fig 8
see also Archicarinae
camaenid snails 200
harvester ants 29–33 & Fig. & Tab. 1, 35–41 &
Tabs 2–4 & Figs 2–3
moth clade: fern associations 245–246 & Fig. 2
resolution of kangaroo phylogeny using DNA
hybridization 310, 323–324
scincid lizards 273–276, 279 Fig. 1
see also
Brisbane, Coffs Harbour, Melbourne, Sydney,
Tasmania and Townsville, North
Carlia and *Lampropholis delicata*
Australia and the South Pacific: patterns in brood
parasitism 262, 271 App.
autapomorphies: Madagascan *Tropidophora* 186,
188–189
authors: patterns in species description 229, 231 Fig.
4, 235–236
autumn spiders *see* *Metellina segmentata*
avoidance learning: genetics of predation 19–23, 26
Azelina: citation as *Ischalis* 250 App.
- badgers (*Meles meles*): dimorphic comparisons 294,
303, 305
- Bagous* (Curculionidae): phytophagous insects 116
- bait: convergence in harvester ant assemblages 29,
33, 38, 44 App.
number occupied 34, 36–37 Tabs 2–3, 44 App.
- BAL (Balloa, Sardinia): *Culex pipiens* sample site
332–334 & Tab. 1 & Fig. 1, 336 Tab. 2
- Balanus improvisus*: epibiont background for isopod
Idotea baltica 48
- Balsaminaceae: insect food-plants 114 Fig. 1, 125
App. 1, 127 App. 2
- Baltic: polymorphism of the isopod *Idotea baltica*
48–49
see also Finland, Orhisaari and Rymättylä 84
- Banksia* spp.: *Lampropholis delicata* experimental areas
277
- Batesian mimicry: butterfly behaviour 159–161, 176
- bats (*Cynopterus* and *Haplonycteris*): gene flow 83
- bees
fit of flowers around their bodies 355–358,
360–361, 364–368, 370–375 & Tab. 6,
378–379
see also bumblebees
patterns in brood parasitism 261–272 & Figs 1–3 &
Tab. 1 & App.
see also halictid bees
- beetles
chrysomelid: insect-plant interactions 248
genetics of predation 17–27 & Tabs 1–3 & Figs 1–2
see also ladybirds and mealworms
variation in morph frequencies 65
see also Coleoptera
- behaviour, feeding (quails): genetics of predation
17–18
- *behavioural ecology: butterfly behaviour and mimicry
159–162
- Benton, M. J., and Spencer, P. S. *Fossil Reptiles of
Great Britain*. (review: F. M. Broadhurst) 385–386
- Berberidaceae: insect food-plants 114 Fig. 1, 119, 125
App. 1, 127 App. 2
- Bettongia penicillata*: resolution of kangaroo phylogeny
using DNA hybridization 315

- Betulaceae
 insect food-plants 110, 112 Tab. 1, 114–115 & Fig. 1, 125 App. 1, 127 App. 2
 larval hosts for *Taeniaria* 250 App.
Betula 247, 250 App.
see also *Alnus*
- Bicyclus anynana* (butterfly pupae): insect prey experiments 19–20 & Fig. 1, 26
- bills (culmens): African whydahs (*Vidua*) 131, 133–134 & Tabs 1–3
- biochemistry: pteridophyte anti-predator strategies 242
- *biodiversity: patterns in species description 225, 234, 236
- Biodiversity and Global Change*, editors: O. T. Solbrig, H. M. van Emden and P. G. W. J. van Oordt.
 (review: Oliver L. Gilbert) 181–182
- biogeographic regions (1–9) 226–235 & Tabs 3–4 & Fig. 3 & 8
see also Afrotropics (4), Australia (8), Eastern Palearctic (6), Indo-Pacific (7), Madagascar (5), Nearctic (1), Neotropics (2), New Zealand (9) and Western Palearctic (3)
- *biogeography: evolutionary genetics of Asian mammals 69–70, 78, 80–81
- BIOSYS-1
 evolutionary genetics of Asian mammals 72
 Madagascar *Tropidophora* 186
- birds
 African whydahs (*Vidua*) 130–131
 colour patterns in a lizard 274–275, 287
see also butcherbirds, currawongs, kookaburras and magpies
 fit of flowers around the bodies of bees 356
see also hummingbirds
 flight display compensations
 male skylarks (*Alauda arvensis*) 138
 male widows (*Euplectes*) 138
 genetics of predation 17–27 & Figs 1–2 & Tabs 1–3
see also quail
 natural selection 130
 passerine: selection of pastry baits 106
 predation of *Heliconius* butterflies 174–175
see also flycatchers and jacamars
 resource limitations 41
 sexual selection acting on females 256 Tab. 1, 258
see also blackbirds, dotterels, flycatchers, fowl, passerines, phalaropes, jacanas, sandpipers, shorebirds, swallows, turkeys and widowbirds
 blackbirds, red-winged: 256 Tab. 1,
 bladderwrack (*Fucus vesiculosus*): habitat choice of adult isopod *Idotea baltica* 45, 48–52, 63–66
- Blechnaceae: lithinine moth hosts 240, 244–245 Tab. 2 & Fig. 2, 250 App.
Blechnum 240, 244–245 Tab. 2 & Fig. 2, 250 App.
Blechnum (Acanthaceae): host to *Siproeta stelenis* (nymphalid butterflies) 240
- 'blue' mimicry rings: butterflies 162, 175
see also *Heliconius doris* and *H. sara* (blue form)
- bluebells: insect food-plants 122
- body size: colour patterns in a lizard 273, 275, 278–288 & Fig. 3
- *body size: whydah (*Vidua*) mass 129, 131–140 & Tabs 1–3 & 6–7
- **Bombus*: fit of flowers around the bodies of bees 355, 365 Figs 2–3, 378
- B. fervidus* 373 Tab. 5 n.
B. impatiens 373 Tab. 5 n.
B. vagans 367–369 & Fig. 6, 371–373 & Tab. 5
- Bonferroni methods
Culex pipiens test significance levels 335
 whydah (*Vidua*) F-ratio comparisons 133, 136
- bootstrap tests
 evolutionary genetics of Asian mammals 78, 80 Fig. 3
 resolution of kangaroo phylogeny using DNA hybridization 309, 312–313, 318–325 & Figs 2–3
- Boraginaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Borneo: *Niviventer* rats 81
- Bowles, M. L. and Whelan, R. J. editors *Restoration of Endangered Species, Conceptual Issues, Planning and Implementation*. (review: Norman W. Moore) 181
- Brachymyrmex*: South American harvester ant assemblages 40
B. giardi 44 App.
B. laevis 44 App.
B. longicornis 44 App.
- bracken
 ant associations 210
 "common brake" (*Pteris*): *Ischalis* host association 250 App.
- Bremen, Germany: patterns in brood parasitism 263–265 & Fig. 2, 271 App.
- Brisbane: *Lampropholis delicata* 276, 279–280 Figs 1–2
- Britain
 insect food-plants 109–127 & Figs 1–8 & Tab. 1 & Apps 1–2
Mustela vison (American mink) 295 Tab. 1, 297–301 & Tabs 1–7, 305
see also England, Ireland and Scotland
see also England
- Buddlejaceae: insect food-plants 110, 114 & Fig. 1, 117, 125 App. 1, 127 App. 2
- buffers: *Culex pipiens* population structure and genetic invasion 331
- bumblebees: fit of flowers around their bodies 355, 357–360, 364, 372–373 & Tab. 5, 377, 379–380
see also bees
- butcherbirds (*Craticus* spp.): lizard predators 287
- Butomaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- butterflies
 behaviour and mimicry 159–180 & Figs 1–4 & Tabs 1–4
 host associations 240
see also lycaenids and nymphalids
 insect quail prey experiments: pupae (*Bicyclus anynana*) 17, 19–20 & Fig. 1, 26
 lycaenid: unpredictable variation in a facultative mutualism 209–223 & Figs 1–4 & Tabs 1–5
see also *Euphilotes enoptes* and *Jalmenus evagoras*
 plasticity 144–145, 153
 speckled wood
Pararge aegeria 143–157 & Figs 1–7
P. xiphia 145
see also females, larvae, males and pupae
- Buxaceae: insect food-plants 114–115 & Fig. 1, 125 App. 1, 127 App. 2

- CAG (Cagliari, Sardinia): *Culex pipiens* sample sites
 CAG1 (tank) 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2, 338
 CAG2 (canal) 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2, 338
- Callidulidae (lepidopteran family): fern associations 241 Tab. 1
 Callidulinae (subfamily) 241 Tab. 1
 Calliduloidea (superfamily) 241 Tab. 1
- calliphorid aphids: insect food-plants 115
- Callitrichaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Callophistria* (noctuid butterflies): pteridophagy 240–241 & Tab. 1
- camaenid snails: Australia 200
- Cambrian spicules: relationships between polyaxonid and monaxonid spicules in Demospongiae 12
- Campanulaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Camponotus*: tending *Euphilotes enoptes* larvae 219
C. vicinus 213–214 & Tab. 1
- Canada: patterns in brood parasitism 262, 270 App
 canals, axial: relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 3–12 & Figs 1–3
- Cannabiaceae: insect food-plants 114 Fig. 1, 119, 125 App. 1, 127 App. 2
- canonical variate analysis (CVA): cranial variation in mink 296, 298–302 & Figs 1–4
- Caprifoliaceae: insect food-plants 112 Tab. 1, 114–115 & Fig. 1, 125 App. 1, 127 App. 2
- capture-mark-recapture studies: colour pattern in a lizard 275, 277, 283–285 & Figs 4–6, 287, 289–290
- Carlia vivax*: juvenile and female lizards with stripes 288
- carnassials: cranial variation in mink 294
- *Carnivora: *Mustela vison* (American mink) 293, 305
- Carnivora: sexual size dimorphism 352
- carotenoids: sexual selection 258
- Caryophyllaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 117, 125 App. 1, 127 App. 2
- Caryophyllidae 120 Fig. 6
- case study: variation in predator preference 103–105 & Tab. 1
- CATMOD: colour pattern polymorphism in a lizard 278
- Cecropia*: Corcovado secondary forest 163
- Celastraceae: insect food-plants 114 Fig. 1, 117, 125 App. 1, 127 App. 2
- Celastrales 113
- censuses
 ants and *Euphilotes enoptes* 212–213
 convergence in harvester ant assemblages 33–34, 44 App.
- Centaurea scabiosa*: insect food-plants 122
- Central America *see* Costa Rica, Mexico and Panama
- Ceractinomorpha (subclass): Demospongiae 12
- Cerambycidae (Coleoptera): phytophagous insects 114
- Ceratophyllaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Cerion*: overdescribed West Indian snails 184
- Cerithium*: Prosobranchia: shell variation 202
- CFR (Arborea, Sardinia): *Culex pipiens* sample site 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2, 338
- chamaephytes: insect food-plants 111, 114–115 & Fig. 2, 117
- character variations, artificial: fit of flowers around the bodies of bees 365–368 & Figs 4–5, 377–380
- Chelaner*: Australian harvester ants 40
- chemical facilitation: lithinine moths on ferns 239–240, 242–244 & Fig. 1, 247–248
- Chenopodiaceae: insect food-plants 112 & Tab. 1, 114 Fig. 1, 116–117, 125 App. 1, 127 App. 2
- CHI (Chilivani, Sardinia): *Culex pipiens* sample site 332–334 & Tab. 1 & Fig. 1, 338
- chi-square tests
 colour pattern polymorphism in a lizard 278
 evolutionary genetics of Asian mammals 73
 polymorphism of the isopod *Idotea baltica* 49, 57, 61
- Chile: harvester ants 31–34 & Fig. 1, 39–40, 44 App. & ns
- China
 Eastern Palearctic (6): south to Yangtze River 226
Niniventer bukit 72, 78, 86 App. 1
- Choreutidae (Lepidoptera): phytophagous insects 111
- chromosomes: sex determination in isopod *Idotea baltica* 48–49
- Chrysomelidae (Lepidoptera)
 insect-plant interactions 248
see also *Ophraella* and *Phyllobrotica* 248
 phytophagous insects 116
- Cirsium acaule*: insect food-plants 122
- Cistaceae: insect food-plants 114–115 & Fig. 1, 125 App. 1, 127 App. 2
- *cladistics: Madagascan *Tropidophora* 183–184, 186, 189–190 & Fig. 2, 200
- cladistics
 cladograms: phylogenetics of Taiwanese small mammals 71 Fig. 1
 lithinine moths on ferns 242–248 & Fig. 1
 resolution of kangaroo phylogeny using DNA hybridization 309, 312, 319
Thylogale-Petrogale-Dendrolagus clade 319
- Cladodia*: spicules in Demospongiae 8
- Cladophora* sp. (green algae): habitat choice of juvenile isopod *Idotea baltica* 45, 48–50, 52–55 & Fig. 2 & Tab. 2, 62–63
- classes, vegetative: patterns in brood parasitism 262–267 & Tab. 1
- *cleptoparasitism: patterns in brood parasitism 261–262, 267
- climates
 patterns in brood parasitism 262–266 & Fig. 3, 270–272 App.
- plasticity in *Pararge aegeria* 143, 145–146
- clines, elevational: evolutionary genetics of Asian mammals 69, 78, 82
- clutches: colour patterns in a lizard 273, 275–276, 281–282 & Tab. 1 & Fig. 3, 288
- co-mimics: *Heliconius* butterflies 159, 161–162, 172–174 & Tab. 4, 177
- Coccinella septempunctata* (seven-spot ladybirds): toxic: genetics of predation 19–21 & Fig. 1, 23, 25–26 & Fig. 2
- Coccinellidae (ladybirds): genetics of predation 17–18
- Coccoidea (scale insects): phytophagous insects 111, 115–116
- Kermisidae 111
- coefficients

- annual precipitation in arid regions: variation 39–40
- cranial variation in mink: principal components (PCs) 297 Tab., 300–301 Tabs 8–9
- fit of flowers around bees: regression 373
- insect food-plants: dissimilarity (Czelkanowski) 111–113
- ornament growth, variation and dimorphism in *Vidua*
- repeatability 132
- variation (CVs) 129, 132, 134–136 & Tabs 2–5
- *coevolution: lithinine moths on ferns 239–243 & Tab. 1, 247
- Coffs Harbour (C): *Lampropholis delicata* 276, 279–282 & Figs 1–3 & Tab. 1
- Cohen's Kappa statistic: cranial variation in mink 296, 302
- Coleoptera: phytophagous insects 114
- see also Cerambycidae and Scolytidae
- *colonization: lithinine moths on ferns 239, 242–244 & Fig. 1, 247–248
- *colour: *Lampropholis delicata* patterns 273–277, 279–282, 285–289
- colour patterns
- American mink see pelt colours
- genetics of predation 17–22
- Japanese quail chick preferences: red vs blue 18, 22
- 'two-spot' ladybirds
- black form ("melanic") 19–21
- red form 17, 19–20
- polymorphisms in a lizard 273–277, 279–282, 286–289
- CATMOD 278
- *colour polymorphism: isopod *Idotea baltica* 45–68 & Figs 1–6 & Tabs 1–5
- see also morphs
- Commelinidae: insect food-plants 113
- communities: harvester ants 29, 34–35, 36–39 & Tabs 2–3 & Fig. 3, 41
- Compositae: insect food-plants 111–112 & Tab. 1, 114 Fig. 1, 117, 125 App. 1, 127 App. 2
- Comptonia*: larval hosts for *Tacparia* 250 App.
- computer programs see BIOSYS-1, CATMOD, CONSENSE, FITCH, GENEPOP, GENSTAT, Hennig86, LINKDOS, MANOVA, PAUP, PHYLIP, SAS and StatGraphics
- conchology: Madagascan *Tropidophora* 183–184, 201
- see also shells
- conclusions
- butterfly behaviour and mimicry 177
- colour pattern polymorphism in a lizard 289–290
- Culex pipiens* population structure and genetic invasion 341
- lithinine moths on ferns 257–248
- polymorphism of the isopod *Idotea baltica* 65–66
- resolution of kangaroo phylogeny using DNA hybridization 325–326
- *condition indicators: sexual selection acting on females 251, 253–254 & Fig. 1, 257
- conifers: insect food-plants 110, 117–118 & Fig. 3
- CONSENSE program: resolution of kangaroo phylogeny using DNA hybridization 312, 320–321 Fig. 2
- consensus trees
- 50\ majority rule: evolutionary genetics of Asian mammals 78, 80 Fig. 3
- Nelson: Madagascan *Tropidophora* 189
- *conservation: Madagascan *Tropidophora* 183
- consistency indices
- evolutionary genetics of Asian mammals 78
- Nelson consensus tree (Madagascan *Tropidophora*) 189
- constraints: convergence in harvester ant assemblages 29, 31, 40–41
- continents
- convergence in harvester ant assemblages 29–31, 33–35, 38–41
- pairwise 29, 35, 40
- three-continents 29, 35–38 & Figs 2–3 & Tabs 2–3, 40
- see also Australia, North and South America
- evolutionary genetics of Asian small mammals 69–71 & Fig. 1, 76, 78, 80–81 & Fig. 3, 85 App. 1, 87 App. 2
- convergence (C values): harvester ant assemblages 29–31, 35–36 & Tab. 2, 39–41
- *convergent evolution: harvester ant assemblages 29–30
- Convolvulaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 119, 125 App. 1, 127 App. 2
- Corcovado, Costa Rica: butterfly behaviour and mimicry in *Heliconius* 162–168 & Fig. 1, 172, 175
- Cornaceae: insect food-plants 109, 112–112 & Tab. 1, 114 Fig. 1, 125 App. 1, 127 App. 2
- Cornales 112
- corpuscles: relationships between polyaxonid and monaxonid spicules in Demospongiae 1–3, 6 Fig. 3, 11
- correlations
- butterfly behaviour and mimicry 159, 161, 166, 168, 170, 172, 174–176
- convergence in harvester ant assemblages 31, 33
- Culex pipiens* population structure and genetic invasion 336
- evolutionary genetics of Asian mammals 70, 78–79 & Fig. 2
- fit of flowers around the bodies of bees 361–363 & Tab. 3, 373, 378, 380
- insect prey experiments 21, 23, 26
- patterns in brood parasitism 263, 267
- Corsica: *Culex pipiens* population structure and genetic invasion 329–331, 335, 337–341 & Tabs 3–5
- see also LUM and POG
- Corylaceae: insect food-plants 110, 112 Tab. 1, 114 Fig. 1, 125 App. 1, 127 App. 2
- *cost of reproduction: sexual selection acting on females 251–259 & Figs 1–2 & Tab. 1
- Costa Rica: butterfly behaviour and mimicry in *Heliconius* 162–165 & Fig. 1, 167, 174–175
- see also Corcovado and Sirena
- Coturnix coturnix japonicus* (Japanese quail): genetics of predation 17–19
- covariances
- morphological traits: whydahs (*Vidua*) 136–138 & Tabs 6–7
- see also ANCOVAs
- Crambe: relationships between polyaxonid and monaxonid spicules in Demospongiae 1–6 & Figs 1–3, 8, 10–12 & Fig. 7

- C. acuta* 3–4 & Fig. 1, 7 & Fig. 4, 10–11 & Fig. 7
C. erecta 2 3
C. taillezi 3 4 & Fig. 1, 5–7 & Figs 2 3, 11
C. tuberosa 1, 3–4 & Fig. 1, 5–6 & Figs 2 4, 9–11 & Figs 6–7
 *craniometrics: variations in mink 293 305 & Tabs 1–9
 see also measurement error and skulls
 Crassulaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
Craticus spp. (butcherbirds): lizard predators 287
 Cruciferae: insect food-plants 112–112 & Tab. 1, 114 Fig. 1, 117, 125 App. 1, 127 App. 2
 Crustacea: *Notonecta* prey
 Amphipoda: *Gammarus pulex* 104–105 & Tab. 1
 Isopoda (*Asellus aquaticus*) 104–105 & Tab. 1
 *crypsis: colour patterns in a lizard 273, 285 286, 288 289
 crypsis: differences in isopod *Idotea baltica* 45 48, 62–65
 *cryptic coloration: butterfly behaviour and mimicry 159
 Cucurbitaceae
 butterfly food plants 161–162, 167
 see also *Gurania* and *Psiguria*
 insect food-plants 114 Fig. 1, 117, 125 App. 1, 127 App. 2
 see also *Bagous*
Culex pipiens (mosquitoes): population structure and genetic invasion 329–343 & Figs 1–2 & Tabs 1–5
C. tritaeniorhynchus 340
 culmens (bills): African whydahs (*Vidua*) 131, 133 & Tab. 1, 137–138 Tabs 6–7
 Cupressaceae
 insect food-plants 114–115 & Fig. 1, 117, 125 App. 1, 127 App. 2
 Thallophaga host 250 App.
 see also *Thuja*
 Curculionidae (Lepidoptera): phytophagous insects 116
 currawongs (*Strepera graculina*): lizard predators 287
Cyamon: relationships between polyaxonid and monaxonid spicules in Demospongiae 11 12
 Cyatheaceae (including Dicksoniaceae): moth associations 242, 244–245 Tab. 2 & Fig. 2, 250 App.
 Cyathea 242, 244–245 Tab. 2 & Fig. 2, 250 App.
 see also *Alsophila* and *Dicksonia*
Cynopterus bats: gene flow 83
 Cyperaceae: insect food-plants 114 Fig. 1, 116, 125 App. 1, 127 App. 2
 Cyperales 120
 Czelkanowski coefficient: principle coordinates analysis (PCA) (insect food-plants) 111–112

D statistics: *Culex pipiens* population structure and genetic invasion 335
Dacelo novaguineae (kookaburras): lizard predators 287
Dalechampia: pollination of blossoms 378
 Danaïnae: butterfly behaviour and mimicry 160
 Daphni, Greece: patterns in brood parasitism 262–267 & Figs 2 3, 271 App.
Daphnia magna (water fleas): *Notonecta* diet 104
 Darwin, C.: sexual ornament and body size 129 130
 data
 colour patterns in a lizard 273–274, 276, 278–281, 285, 289
 convergence in harvester ant assemblages 31, 33, 39
 ecological: Madagascan *Tropidophora* 190, 192–193 Tab. 2, 199 200, 202
 fit of flowers around the bodies of bees 357, 359 361, 363 364, 371, 377, 379 380
 insects
 food-plants 109–111, 123
 major 'data base' 234
 ornament growth, variation and dimorphism in *Vidua* 129–133 & Tab. 1
 patterns
 brood parasitism 263
 species description 226 227, 235
 daylengths: plasticity in *Pararge aegeria* 143 144, 147 154 & Figs 2 7
Delphinium nelsoni: pollination of albino flowers 380
 *demosponges: relationships between polyaxonid and monaxonid spicules 1–15 & Figs 1 7
 dendrograms
 UPGMA: evolutionary genetics of Asian mammals 77–79 & Fig. 2, 81–82
 see also trees
Dendrolagus: resolution of kangaroo phylogeny using DNA hybridization 310–311, 315, 319, 324 326
D. dorianus 315
D. inustus 315
D. matschiei (tree kangaroos) 309, 314–316 & Tabs 1–2 & Fig. 1, 320–321 Fig. 2
Dendrolagus-Thylogale Petrogale association 309
 Dennstaedtiaceae: moth associations 242, 245 246 & Fig. 2, 248
Dennstaedtia (*Homochloides*): *Ischalia* larval hosts 250 App.
 see also *Histiopteris*, *Paesia* and *Pteridium*
 *deserts: convergence in harvester ant assemblages 29–31, 39–41
 desmas: relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 3–4 & Fig. 1, 7–11 & Figs 4 & 7
 desmoids, asterose: relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 3, 10 11
 development: plasticity in *Pararge aegeria* 143–148
 pathways 143–144, 146–151 & Figs 1–5
 times 143–144, 146 154 & Figs 1–7
 *diapause: plasticity in *Pararge aegeria* 143 151 & Fig. 2, 153
Dicksonia (Cyatheaceae: tree ferns): moth associations 242, 244 245 Tab. 2 & Fig. 2, 250 App.
 Dicksoniaceae 245 Fig. 2, 250 App.
 Dicotyledonae
 insect food-plants 113, 116
 Magnoliopsida: larval hosts for *Lithini* 250 App.
 see also Betulaceae, Myricaceae, Salicaceae and Saxifragaceae
 Dilleniidae: insect food-plants 119–120 & Fig. 6
 *dimorphic niches: polymorphism of the isopod *Idotea baltica* 45, 64
 *dimorphism: sexual selection acting on females 251 252, 257 258
 dimorphism
 cranial variation in mink 293 295, 298–301 & Figs 2 4 & Tabs 4–7, 304–305

- Lampropholis delicata* 273
Metellina segmentata 345–354 & Figs 1–3 & Tab. 1
 whydahs (*Vidua*) 129, 132–133 & Tab. 1, 136, 138–139
Dione juno (butterflies): 'orange' mimicry rings 165
 Dioscoraceae: insect food-plants 120–121
 Dipsacaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
 Diptera: phytophagous insects 116–117
Discorhabdella: relationships between polyaxonid and monaxonid spicules in Demospongiae 1–6 & Figs 1–2, 8 Fig. 5, 11–12
D. hindei 2, 4–5 & Figs 1–2
D. incrustans 2–5 & Figs 1–2, 7–9 & Figs 5–6
D. tuberosocapitata 2, 4–5, 8 Fig. 5
 discussions
 butterfly behaviour and mimicry 174–176
 colour patterns in a lizard 285–290
 convergence in harvester ant assemblages 39–41
 cranial variation in mink 302–306
Culex pipiens population structure and genetic invasion 339–341
 evolutionary genetics of Asian mammals 78, 80–83
 fit of flowers around the bodies of bees 377–381
 genetics of predation 17, 24–27 & Fig. 2
 insect food-plants 109, 121–123
 lithinine moths on ferns 246–247
 Madagascar *Tropidophora* 200–202
 ornament growth, variation and dimorphism in *Vidua* 136, 138–140
 patterns
 brood parasitism 265–267 & Fig. 3
 species description 234–236
 plasticity in *Pararge aegeria* 143, 155
 polymorphism of the isopod *Idotea baltica* 61–66
 relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 8–12 & Figs 6–7
 resolution of kangaroo phylogeny using DNA hybridization 319–325 & Figs 2–3
 sexual dimorphism in *Metellina segmentata* 350–352
 sexual selection acting on females 257–259
 unpredictable variation in a facultative mutualism 216–221 & Figs 2–4 & Tabs 2–5
 variation in predator preference 105–106
 Dismophiinae: butterfly behaviour and mimicry 160
 distributions
 Taiwanese small mammals 69–71, 81
 variation in predator preference
 bimodal 97–101 & Figs 3–4
 normal 101–103 & Figs 5–6
 diversification: genetics of Taiwanese small mammals 69, 71, 73–74, 77, 80, 82
 *DNA hybridization: resolution of kangaroo phylogeny 310–311, 319, 323–326
 DNA sequences: evolutionary genetics of Asian mammals 80–81
 Dolichoderinae
 harvester ant assemblages 40
 tending *Euphilotes enoptes* larvae 213 Tab. 1
Tapinoma sessile 213 Tab. 1
 *domestication: American mink 293–294, 303–305
Dorcopsulus: resolution of kangaroo phylogeny using DNA hybridization 309, 313, 319, 323–324, 326
D. vanheurni (forest wallaby): New Guinea 309, 311, 313–316 & Tabs 1–2 & Fig. 1, 318 Tabs 3–4, 320–321 Fig. 2
Dorymymex: South American harvester ant assemblages 40
D. ensifer 44 App.
D. exsanguis 44 App.
 dotterels: sexual selection acting on females 256 Tab. 1,
 *drift: *Culex pipiens* population structure and genetic invasion 329–331, 335–336, 339–340
Drosera tracyi: pollination of trimmed flowers 380
Dryadula phaeusa (butterflies): 'orange' mimicry rings 165
Dryas iulia (butterflies): 'orange' mimicry rings 165
 Dryopteridaceae (*Polystichum*): moth associations 242, 244–245 Tab. 2 & Fig. 2, 250 App.
 see also *Dryopteris*, *Polystichum* and *Rumobra*
Dryopteris (*Gueneria* and *Petrophora*): *Ischalis* host citations 250 App.
D. pennigera (*Pneumaopteris*) 250 App.
 Dundrum, Co. Down: sexual dimorphism in *Metellina segmentata* 346–347
 Eastern Palaearctic: biogeographic region (6) 226, 228–231 Tabs 1–2 & Fig. 3, 231 Fig. 8
 see also China
 ecology
 colour patterns in a lizard 276, 285–287
 Madagascar *Tropidophora* 190, 192–193 Tab. 2, 199–200, 202
 *ecology, Mediterranean: patterns in brood parasitism 261, 266–267
 *ecosystems: patterns in brood parasitism 261–262, 267
 see also arid and subarid
 Ecuador: *Heliconius cydno* 176
 eelpout (*Zoarces viviparus*): isopod *Idotea baltica* experiments 50–51, 59 & Fig. 4, 64
 Elaeagnaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
 Elasmopeidole: South American harvester ant assemblages 39–40
E. subaberrans 44 App.
Electra crustulenta: epibiont background for isopod *Idotea baltica* 48
 electrophoreses
Culex pipiens population structure and genetic invasion 330–331
 evolutionary genetics of Asian mammals 69, 71–72, 85–86 App. 1
 Madagascar *Tropidophora* 186–193 & Tabs 1–2 & Figs 3–4, 197 Fig. 8, 205–208 App.
 *elevational cline: evolutionary genetics of Asian mammals 69–71, 73, 76–78, 81–82
 Ellinikón nr Daphni, Greece: patterns in brood parasitism 266 Fig. 3
 Empetraceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
 *endemism: variation: Madagascar *Tropidophora* 183
 England
 American mink farms see Kitson and Peregrine
Pararge aegeria (speckled wood butterflies) 143–145, 147 & Fig. 1, 150 & Fig. 5, 152–153
 Ennominae
 looper moths: fern associations 241 & Tab. 1

- patterns in species description 227–228 & Figs 1–2, 232–233 & Tabs 3–4
- Entomopteryx* (*Mesophlebion*): *Lithinini* larval hosts 250
- App.
- environments
- convergence in harvester ant assemblages 29, 39, 41
 - differential predation affects colour polymorphism of isopod *Idotea baltica* 45–46, 62
 - plasticity in *Pararge aegeria* 143–144, 146
- enzymes
- Culex pipiens* population structure and genetic invasion 331
 - see also AAT, ACO, FUM, GPD, GPI, HK, IDH, MDH, ME, ODH, PGD and PGM
 - evolutionary genetics of Asian mammals 72
 - see also allo- and isoenzymes
- epibionts: background for isopod *Idotea baltica* 48
- see also *Balanus improvisus*, *Electra crustulenta*, *Mytilus edulis* and *Theodoxus fluviatilis*
- Equisetaceae (horsetails): insect food-plants 114 Fig. 1, 116–117, 125 App. 1, 127 App. 2
- Ericaceae: insect food-plants 112 Tab. 1, 114–115 & Fig. 1, 125 App. 1
- Eriogonum* (Polygonaceae): *Euphilotes enoptes* host 211, 220 Fig. 4
- E. compositum* 209, 211–212
- Est-1*: Madagascan *Tropidophora* (allozyme data) 186, 189 Fig. 2, 205–206 App.
- Est-2*: Madagascan *Tropidophora* (allozyme data) 186, 189 Fig. 2, 205–206 App.
- esterases: population structure and genetic invasion 330
- see also A2B2
- Eucalyptus* spp. (gum trees): *Lampropholis delicata* experimental areas 277
- Euonymus*: insect food-plants
- E. europaeus* 116
 - E. japonicus* 116
- Euphilotes enoptes* larvae: unpredictable variation in a facultative mutualism 209–221 & Tabs 1–5 & Figs 1–4
- Euphorbiaceae: insect food-plants 109, 112–114 & Tab. 1 & Fig. 1, 125 App. 1, 127 App. 2
- Euphorbiales 113
- Euplectes* (widows)
- male flight display compensations 138
 - ornament length: *E. prognus* 139
- Eurasia: evolutionary genetics of Asian mammals 70, 78
- see also Serbia
- Europe
- American mink 294, 305
 - see also Britain, Ireland and Norway
 - Culex pipiens*: see also Corsica, France and Sardinia
 - Metellina segmentata*: northern Europe 345–346
 - Pararge aegeria* (speckled wood butterflies) 145
 - see also Baltic, Britain, England, Finland, Germany, Greece, Madeira, Serbia, Spain, Sweden and Switzerland
 - species descriptions: Western Palaearctic 226
- Eurypon coronula*: relationships between polyaxonid and monaxonid spicules in Demospongiae 3, 6
- Eutheria, herbivorous: resolution of kangaroo phylogeny using DNA hybridization 310
- evergreens: insect food-plants 109, 115–116
- *evolution: relationships between polyaxonid and monaxonid spicules in Demospongiae 1–2, 12
 - *evolution: sexual dimorphism in *Metellina segmentata* 345
- evolution
- convergence in harvester ant assemblages 29–31, 40–41
 - cranial variation in mink 294
 - fit of flowers around the bodies of bees 356, 358, 372, 380
 - genetics
 - Asian mammals 71, 78, 80–81
 - predation 17–18
 - insect food-plants, life form and taxonomy 109–110, 112, 121–123
 - ornament growth, variation and dimorphism in *Vidua* 130
- Exodictia*: relationships between polyaxonid and monaxonid spicules in Demospongiae 8
- experiments
- case study of variation in predator preference 94, 104–105 & Tab. 1
 - fit of flowers around the bodies of bees 355, 357, 359–365, 368–373 & Figs 6–7, 377–380
 - see also character variations
 - genetics of predation
 - F1 and F2 generations 17–23 & Fig. 1 & Tab. 1
 - subsidiary tests (F3 generation) 17–18, 22–23
 - painting: colour patterns in a lizard 273–275, 277–278, 283–289 & Figs 4–6 & Tab. 2
 - plasticity in *Pararge aegeria* 143, 146–147, 150–151
 - polymorphism of the isopod *Idotea baltica* 47–52 & Tab. 1, 55–65 & Tab. 5 & Figs 4–6
- F statistics
- Culex pipiens* migration estimates 335–340 & Tabs 2 & 5
 - ornament growth, variation and dimorphism in *Vidua* 132–134, 136
- *F-statistics: evolutionary genetics of Asian mammals 69, 73, 76–77 & Tab. 3, 82
- see also fixation index
- facilitation, chemical: lithinine moths on ferns 239–240, 242–244 & Fig. 1, 247–248
- Fagaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 125 App. 1, 127 App. 2
- Fagales 109, 117
- 'fallacy of averages': variation in predator preference 94
- families: insect food-plants 109–110
- farms, American mink 294, 297
- see also Heriot, Kitson and Peregrine
- 'fearfulness': insect prey experiments 17, 19, 21–23, 26–27
- fecundity: sexual selection acting on females 251–257 & Figs 1–2
- feeding behaviour (Japanese quail): genetics of predation 17–18
- female phase: fit of flowers around the bodies of bees 358–360 & Fig. 1, 379
- females
- colour patterns in a lizard 273–290 & Figs 1–6 & Tabs 1–2
 - gravid 273, 281–282, 286, 288–290
 - cranial variation in mink 293–295 & Tab. 1,

- 297–298 & Tab. 3, 300–302 & Figs 3–4 & Tabs 6 & 8, 304–305
- Heliconius*: mating behaviour 166–167, 176
- insect prey experiments: Japanese quail 19–20, 23
- Madagascan *Tropidophora* 188, 190–193 & Figs 3–4 & Tab. 2
- ornament growth variation and dimorphism:
- African whydahs (*Vidua*) 129–133, 136, 140
- Pararge aegeria* (speckled wood butterflies) 146–147 & Fig. 1, 149 Figs 3–4, 152, 154 Figs 6–7
- patterns in brood parasitism 262
- polymorphism of the isopod *Idotea baltica* 45, 47–65 & Tabs 1 & 3 & 5 & Figs 1 & 3–6
- neofemales 48–49
- sexual dimorphism in *Metellina segmentata* 345–352 & Tab. 1 & Fig. 2
- sexual selection processes 251–258 & Figs 1–2
- variation in predator preference: waterboatmen (*Notonecta*) 93–107 & Figs 1–6 & Tab. 1
- feral populations: cranial variation in mink 293–295 & Tab. 1, 298–302 & Tabs 5–6, 304–305
- ferns
- insect food-plants 110, 116
- see also Polypodiaceae
- lithinine moths 239–250 & Figs 1–2 & Tabs 1–2 & App
- ferrets (*M. putorius furo*): domestication 305
- *Filicopsida: lithinine interactions 239–242 & Tab. 1, 246–247, 250 App.
- finches, African: whydahs (*Vidua*) 129–130
- Finland: isopod *Idotea baltica* 49–50
- see also Gulf of Bothnia 50
- fireweed aphids and ants 210
- fish
- predation: polymorphism of the isopod *Idotea baltica* 47, 50, 62
- sex-linked colour polymorphism 274
- *Fisher process: sexual selection acting on females 251–254 & Fig. 1
- Fisher's tests: *Culex pipiens* population structure and genetic invasion
- combined probability 332–333 & Tab. 1, 335–337 & Tabs 2–3
- exact test 334–335, 337 Tab. 3
- FITCH program: resolution of kangaroo phylogeny using DNA hybridization 312–313, 315, 318–321 & Fig. 2, 324–325
- fitness
- colour patterns in a lizard 273–274, 289
- fit of flowers around the bodies of bees 357, 366, 378–379
- sexual selection acting on females 251–252, 255, 257
- fixation index: evolutionary genetics of Asian mammals 73
- fleas, water (*Daphnia magna*): *Notonecta* diet 104
- flight heights: butterfly behaviour and mimicry 159, 161–163 & Tab. 1, 165–168 & Fig. 2, 170–171 & Tab. 3, 174–175, 177
- *floral morphology: fit of flowers around the bodies of bees 355–373 & Figs 1 & 4–8 & Tabs 2–4, 375–380 & Tabs 7–8
- flycatchers
- pied: sexual selection acting on females 256 Tab. 1
- tyrannid: *Heliconius* predation 174
- Folititpa acuatasee* *Crambe acutata* 11
- food plants: butterfly behaviour and mimicry 161–162
- see also Cucurbitaceae and Passifloraceae
- foraging modes: harvester ant assemblages 29, 33
- distances 29, 34, 36–38 & Tabs 2–3 & Fig. 4
- scores 29, 34–38 & Tabs 2–3
- Forelius*: South American harvester ant assemblages 40
- F. grandis* 44 App.
- Formica* (genus): tending *Euphilotes enoptes* larvae 215
- F. argentea* 213 & Tab. 1
- F. cf. integroides* 213 & Tab. 1
- F. densiventris* 213 Tab. 1
- F. haemorrhoidalis* 213 Tab. 1
- F. manni* 213 Tab. 1
- F. neogagates* 213 & Tab. 1
- F. oreas* 213 Tab. 1
- F. species* 213–214 & Tab. 1, 219
- F. subpolita* 213 & Tab. 1
- Formicinae
- convergence in harvester ant assemblages 40
- unpredictable variation in a facultative mutualism 213 Tab. 1
- Formicidae 209
- see also Camponotus, *Formica* and *Lasius*
- fossil records
- Macropodinae and collateral subfamilies 310
- relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 3, 7–11 & Fig. 4
- Fossil Reptiles of Great Britain, by M. J. Benton and P. S. Spencer. (review: F. M. Broadhurst) 385–386
- fowl, mallee: sexual selection acting on females 256 Tab. 1,
- France: *Culex pipiens* population structure and genetic invasion 340–341
- Freehold, Greene County: fit of flowers around the bodies of bees 361–362 & Tabs 1–2, 373–376 Tabs 5–8
- *frequency-dependent selection: variation in predator preference 93–107 & Figs 1–6 & Tab. 1
- Fucus*: habitat choice of adult isopod *Idotea baltica* 48, 52–55 & Fig. 2 & Tab. 2
- F. vesiculosus* (bladderwrack) 45, 48–52, 63–64
- FUM (fumerase): *Culex pipiens* population structure and genetic invasion 331
- Fum* locus 333 Tab. 1, 336–337 & Tab. 3
- Fumariaceae: insect food-plants 114 Fig. 1, 119, 125 App. 1, 127 App. 2
- Fungi: insect food-plants 110–111, 113–118 & Figs 1–3, 123, 125 App. 1, 127 App. 2
- G6pd*: Madagascan *Tropidophora* (allozyme data) 186, 205–206 App.
- G-tests
- colour pattern polymorphism in a lizard 279–280, 282–283, 385
- rates of ant attendance: larvae of *Euphilotes enoptes* 213, 215–216 Fig. 1 & Tab. 2, 218 Tabs 4–5
- Galbutia* spp (jacamars): *Heliconius* predation 175
- G. ruficauda* 165
- gall-midge host plant evolution 122
- gammarids
- Gammarus pulex* (Crustacea: Amphipoda) *Notonecta* prey 104–105 & Tab. 1
- predator food in isopod *Idotea baltica* experiments 50
- *Gastropoda: Madagascan *Tropidophora* 183

- gene flow: colour patterns in a lizard 273
- *gene flow: *Culex pipiens* population structure and genetic invasion 329–331, 339–341
- *gene flow: evolutionary genetics of Asian mammals 69, 71, 73, 77 7 Tab. 4, 82 83
see also *Cynopterus* and *Haplonycteris*
- GENEPOP software: *Culex pipiens* computations 333–334
- generations (Japanese quail): insect prey experiments 17–21
F1 19 23 & Tab. 1, 26
F2 19 23 & Tab. 1, 26
F3 (subsidiary test) 17–18, 22, 25 & Fig. 2
- genes
resistance (A2–B2): *Culex pipiens* population structure and genetic invasion 329–330, 340–341
sex determination in isopod *Idotea baltica* 48–49
sexual selection acting on females 254, 257
'good genes' 252, 254–255, 257
- *genetic differentiation: *Culex pipiens* population structure 329, 334–335, 337 340 & Tabs 4–5 & Fig. 2
- genetics
distances
evolutionary genetics of Asian mammals 72, 75–76 & Tab. 2, 81
see also Nei's and Roger's distances
influences: cranial variation in mink 293 295, 303–306
population structures: evolution of Asian mammals 69 71, 73, 76 77
variation: rate of recruitment of novel insect prey into quail diet 17–18
- Genistaceae: insect food-plants 117
- *genitalia: male Madagascan *Tropidophora* 183 184, 188, 195 & Fig. 6, 201
see also penes
- GENSTAT (statistical package): insect food-plants 113
- Gentianaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Geodia*: relationships between polyaxonid and monaxonid spicules in Demospongiae 2, 9
- *Geometridae: lithinine moths on ferns 239, 241 & Tab. 1, 244, 246–247, 250 App.
Geometroidea (superfamily) 241 Tab. 1
- Geometridae: patterns in species description 225–237 & Figs 1–8 & Tabs 1–4
Geometrinae 227 228 & Figs 1–2 & Tab. 1, 232–233 & Tabs 3–4
see also Ennominae, Geometrinae, Laurentinae, Oenochrominae and Sterrhinae
- geophytes (bulbs etc.): insect food-plants 111, 115–117 & Fig. 2, 122
- GER (Geremeas, Sardinia): *Culex pipiens* sample site 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2
- Geraniaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Geraniales 113
- Germany: patterns in brood parasitism 263–265 & Fig. 2, 271 App.
see also Bremen
- Gilbert, J., Danielpool, D. L. and Stanford, J. A.
editors: *Groundwater Ecology* (review: A. A. Savage) 385
- Gleicheniales (ferns): *Ischalis* associations 244–246 & Tab. 2 & Fig. 2, 250 App.
- Gelechioidea (superfamily) 241 Tab. 1
- glucose-6-phosphate (*Gd*, E.C. 1.1.1.49) 72
Niviventer 88 App. 2
- glucose-phosphate-isomerase (*Gpi*, E.C. 5.3.1.9) 72
Apodemus 87 App. 2
Microtus 89 App. 2
Niviventer 88 App. 2
- glutamate oxalate transaminase (*Got-1*, *Got-2*, E.C. 2.6.1.1.) 72
Got-1, *Got-2*: *Microtus* 89 App. 2
Got-2: *Niviventer* 88 App. 2
- glyceraldehyde-3 phosphate-dehydrogenase (*Gapdh*, E.C. 1.2.1.12) 72
- glycerol-3 phosphate dehydrogenase (*Gpd*, E.C. 1.1.1.8) 72
Apodemus 87 App. 2
Microtus 72
Niviventer 88 App. 2
- Gonophylla* on *Aspidium* (*Polystichum*): *Ischalis* associations 250 App.
- gophers, pocket (*Thomomys bottae*)
- gene flow 82
- genetic structure 70
- GPD (glycerol-3 phosphate dehydrogenase): *Culex pipiens* population structure and genetic invasion 331
Gpd locus 332 Tab. 1, 336 338 & Tabs 3–4
- GPI (glucose-6 phosphate isomerase)
Culex pipiens population structure and genetic invasion 331
Gpi locus 332 Tab. 1, 337 338 Tabs 3 4
Madagascan *Tropidophora* (allozyme data): *Gpi* 186, 205 206 App.
- Gramineae
host grasses (*Poa annua*): *Pararge aegeria* (speckled wood butterflies) 145–147
insect food-plants 112 Tab. 1, 114 Fig. 1, 116, 125 App. 1, 127 App. 2
- Graphium*: host-plant relationships 247–248
- grass trees (*Xanthorrhoea* spp.): *Lampropholis delicata* experimental areas 277
- Greece: patterns in brood parasitism 262 266 & Figs 2–3, 271 App.
see also Daphni and Ellinik\ 'a2n
- *gregarious roosting: *Heliconius* butterfly behaviour and mimicry 159–163, 166, 168–177 & Tabs 2–4 & Fig. 3
- Grossulariaceae: insect food-plants 112–114 & Tab. 1 & Fig. 1, 125 App. 1, 127 App. 2
- Groundwater Ecology*, editors: J. Gilbert, D. L. Danielpool and J. A. Stanford. (A. A. Savage) 385
- group principal component analysis (MGPCA): cranial variation in mink 296
- *growth: plasticity in *Pararge aegeria* 143
- Gueneria* (*Dryopteris*): *Ischalis* host citations 250 App.
- Gulf of Bothnia, Finland: isopod *Idotea baltica* 50
- Gurania* spp., Cucurbitaceae: butterfly food plants 161–162
- Gymnorhina tibicen* (Australian magpies): lizard predators 287
- Gymnospermae
insect food-plants 114–115

- larval hosts for Lithiniinae 250 App.
gynoeceum: fit of flowers around the bodies of bees 358
see also stigma
- *habitat choice: polymorphism of the isopod *Idotea baltica* 45 48, 50–53, 62–63, 65–66
microhabitats 46, 48, 52, 62, 65
- habitats
butterfly behaviour and mimicry 159–163, 167 169 & Tab. 2, 172, 174–178 & Fig. 4
colour patterns in a lizard 273–275, 277, 289
harvester ant assemblages 30–31, 33, 39
insect food-plants 109–110, 113, 116–117, 120–123
see also microhabitats
- Hadromerida: spicules in Demospongiae 1, 11–12
- Hainan, Southeast Asia: *Niviventer* rats 81
- Haldane's exact test: *Culex pipiens* computations 331, 334
- halictid bees: patterns in brood parasitism 266
- halophytes: insect food-plants 116
- Haloragaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Hamamelidae: insect food-plants 117–118 & Fig. 4
- *handicap: sexual selection acting on females 251, 254
- Haplonycteris* bats: gene flow 83
- Hardy-Weinberg expectations: *Culex pipiens* population structure and genetic invasion 329, 331–333 & Tab. 1, 336–337 & Tab. 2
- harvester ants *see* ants, harvester
- hawkmoth pollination: *Viola cazorlensis* 379
- Helianthus grosseserratus*: pollination of trimmed flowers 380
- Heliconiinae: butterfly behaviour and mimicry 160–162, 166, 174, 176–177
acraeoid and danoid Heliconiidae: Bates, H. W. 160
- Heliconius*: butterfly behaviour and mimicry 159–180 & Figs 1–4 & Tabs 1–4
groups
erato 166, 172–174, 176–177
melponeme 167, 172–174 & Tab. 4, 176–177
H. charitonia (non-mimetic) 164–166 & Tab. 1, 168–173 & Figs 2–3 (cha) & Tabs 2–4, 175
H. cydno 176–177
H. doris 164–167 & Fig. 1, 168–171 & Figs 2–3 (dor) & Tabs 2–3, 173 Tab. 4, 176
H. erato 163–166 & Fig. 1, 168–177 & Figs 2–3 (era) & Tabs 2 4
H. ethilla 165
H. hecale 164–165 & Fig. 1, 167–173 & Figs 2–3 (hec) & Tabs 2–4, 177
H. heurysoni 164–167 & Fig. 1, 168–173 & Figs 2 3 (hew) & Tabs 2–4
H. ismenius 164–165 & Fig. 1, 167–173 & Figs 2–3 (ism) & Tabs 2–4, 177
H. melpomene 164–165 & Fig. 1, 167–173 & Figs 2–3 (mel) & Tabs 2–4, 176–177
H. numata 160, 176
H. pachinus 164–165 & Fig. 1, 167–173 & Figs 2–3 (pac) & Tabs 2–4, 176–177
H. sara 164–166 & Fig. 1, 169–173 & Tabs 2–4 & Fig. 3
order: *sara-heurysoni-pachinus-melpomene-erato-hecale-ismenius* 169
- 'silvaniforms' 160, 165
- helophytes (marsh plants): insect food-plants 111, 113, 115–116 & Fig. 2
- hemipterophytes: insect food-plants 111–117 & Fig. 2
- Hemiptera
insect food-plants 117
variation in predator preference: *Notonecta* 104
- Hennig86: Madagascan *Tropidophora* 186, 189 Fig. 2
- herbivores
eutherian: resolution of kangaroo phylogeny using DNA hybridization 310
lithinine moths on ferns 240–241 & Tab. 1
- herbs: insect food-plants 109, 116–117, 119, 121–123
- Heriot Farm, Scotland: American mink 299 Fig. 2 & Tab. 4
- Heterodisca*: fern associations 245 Fig. 2
- heterogeneity (G) tests: isopod *Idotea baltica* colour morph changes 52–55 & Tab. 2, 57 & Tab. 3, 59 & Fig. 4
- Heteroptera: phytophagous insects 122
- heterozygosity: evolutionary genetics of Asian mammals 72–75 & Tab. 1, 81–83
- Hippocastaneaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 125 App. 1, 127 App. 2
- Hippuridae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- Hirundo rustica* (barn swallows)
male European 258
male North American 258
- Histiogaster*: moth associations 245–246 & Fig. 2, 250 App.
- history
convergence in harvester ant assemblages 29–31, 34–41 & Tab. 2
genetics of Asian mammals
evolutionary 78, 80–81
geological 70–71, 81
- HK (hexokinase): *Culex pipiens* population structure and genetic invasion 331
Hk1 Hk2 pair 336
Hk-1 locus 336
Hk-2 locus 332 Tab. 1, 334, 336–337 & Tab. 2
- holotypes: *Crambe* and *Discorhaddella* 2–3
- Homochlodes* (*Dennstaedtia* and *Pteridium*): *Ischalia* larval hosts 250 App.
- homology: relationships between polyaxonid and monaxonid spicules in Demospongiae 1–2, 8–9, 10–12
- Homosclerophorida: spicules in Demospongiae 12
- horsetails (Equisetaceae): insect food-plants 114 Fig. 1, 116–117, 125 App. 1
- *host associations: lithinine moths on ferns 239–248 & Tab. 1
- hosts
patterns in brood parasitism 261–262, 266
plants
butterfly behaviour and mimicry in *Heliconius* 161–162, 166–167
see also Cucurbitaceae, *Lantana* and Passifloraceae
grasses (*Poa annua*): *Pararge aegeria* (speckled wood butterflies) 145–147
- human-chimpanzee-gorilla trichotomy 311
- hummingbirds: fit of flowers around their bodies 357–358, 364, 378–379

- Hydrangeaceae: insect food-plants 112
- Hydrocharitaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- hydrophytes (water plants): insect food-plants 109, 111, 113, 115–116 & Fig. 2
- Hymenophyllales (ferns): *Ischalis* associations 244–246 & Tab. 2 & Fig. 2, 250 App.
see also Blechnaceae, Cyatheaceae, Dennstaedtiaceae, Dryopteraceae and Thelypteridaceae
- Hymenoptera: patterns in brood parasitism 261–262
- Hymeraphia spinispinosa*: spicules in Demospongiae 11
- Hypericaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 125 App. 1, 127 App. 2
- Hypochryops theon* (lycaenid butterflies): fern-feeding 240
- hypotheses
- colour patterns in a lizard 273, 275, 279–280, 282, 286 289
 - evolutionary genetics of Asian mammals 69, 71 & Fig. 1, 80
 - null hypothesis 73, 76 Tab. 3 & n.
 - fit of flowers around the bodies of bees 361, 365–367 & Figs 4–5
 - lithinine moths on ferns 239–240, 242, 246–248
 - mimetic diversity in *Heliconius* 176–177
 - sexual dimorphism in *Metellina segmentata* 346
- Iberian mint: variations in pollinators 210
- Icaricia acmon* larvae and ants 219
- IDH (isocitrate dehydrogenase): *Culex pipiens*
 population structure and genetic invasion 331
Idh-1 locus 332 Tab. 1, 336–338 & Tabs 3–4
Idh-2 locus 332 Tab. 1, 337–338 Tabs 3–4
- Idiodes* (loopers moths): Indo-Australian region 241, 250 App.
- Idotea baltica* (Pallas): differential predation on sexes
 effects colour polymorphism 45 68 Figs 1–6 & Tabs 1–5
 adults 45, 47–50, 54–56, 62–63, 65
 juveniles 45, 47–55 & Figs 1–2 & Tab. 2, 57, 62–64
 phenotypes
albafusca 49, 65
flavafusca 49
maculata 49, 65
 sex-reversals
I. b. baltica 49
I. b. basteri 49
I. b. tricuspidata 49
see also females, males and morphs
- Impatiens*: fit of flowers around the bodies of bees 358, 362, 365, 371–372, 376 Tab. 8, 378
I. capensis 355, 357–373 & Figs 1 & 3 & 5 & 7–8 & Tabs 1–5, 377–378
I. pallida 355, 357–373 & Figs 1–2 & 4 & 6 & Tabs 1–5, 377–378
- incursions: Asian mammals into Taiwan 69, 71 & Fig. 1, 80
- indicators: sexual selection acting on females 251–254 & Fig. 1, 257–258
- indices
- convergence in harvester ant assemblages 34–35
 - cranial variation in mink 298
 - evolutionary genetics of Asian mammals
 consistency 78
 fixation 73
 harvester ant assemblages: Shannon diversity 34
 insect prey experiments: 'fearfulness' 19, 21–22
 Madagascan *Tropidophora* *see* consistency and retentive
 Morisita's (I_m): assessment larval aggregation 213, 216, 218 Tab. 3
 Ohta: population structure and genetic invasion 335–336
 variations in predator preferences 96, 105 Tab. 1
see also Manly's and Murdoch's indices
 - *individual-based models: variation in predator preference 93–107 & Figs 1–6 & t. 1
- Indo-Australian region: *Idiodes* (looper moths) 241
- Indo-Pacific: biogeographic region (7) 227–229 & Tabs 1–2 & Fig. 3, 234 lf. 8
- ingroups: resolution of kangaroo phylogeny using DNA hybridization 309, 311, 319, 322–323
- insecticides: *Culex pipiens* population structure and genetic invasion 329–330, 338, 340–341
- insects
- adaptations 143–144
 - fit of flowers around the bodies of bees 357
 - food-plants, life forms and taxonomy 109–127 & Figs 1–8 & Tab. 1 & Apps 1–2
 scale insects (Coccoidea) 111, 115–116
see also aphids, mites and Lepidoptera
 - novel prey: quail experiments 17–19
 - plant associations: lithinine moths on ferns 239–244 & Fig. 1, 246–248
 sex-linked colour polymorphism 274
 - *insects (Geometridae): patterns in species description 225, 236
- instars: *Euphilotes enoptes* 209, 212–216 & Fig. 1 & Tab. 2, 219 220 & Fig. 4
- introductions
- butterfly behaviour and mimicry 160–162
 - colour patterns in a lizard 273–275
 - convergence in harvester ant assemblages 29–31
 - cranial variation in mink 293–295
 - Culex pipiens* population structure and genetic invasion 329–330
 - evolutionary genetics of Asian mammals 69–71
 - fit of flowers around the bodies of bees 356 357
 - genetics of predation 17–18
 - insect food-plants, life form and taxonomy 109–110
 - isopod *Idotea baltica* 45 48
 - lithinine moths on ferns 239–240
 - Madagascan *Tropidophora* 184–186
 - ornament growth, variation and dimorphism in *Vidua* 129–131
- patterns
- brood parasitism 261–262
 - species description 225–226
- plasticity in *Pararge aegeria* 143–145
- relationships between polyaxonid and monaxonid spicules in Demospongiae 1–2
- resolution of kangaroo phylogeny using DNA hybridization 310–311
- sexual dimorphism in *Metellina segmentata* 345 346
- sexual selection acting on females 251–252
- unpredictable variation in a facultative mutualism 210–211
- variation in predator preference 93–95 & Fig. 1
- Ipomopsis*: fit of flowers around the bodies of bees 380
- I. aggregata* 379

- Ireland
Mustela vison (American mink) 295 Tab. 1, 300–301 & Fig. 4 & Tab. 7
 sexual dimorphism in *Metellina segmentata* 345–346
see also Dundrum
- Iridaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- **Ischalis* (lithinine moths): pteridophagy 239, 244–246 & Tab. 2 & Fig. 2, 248, 250 App.
I. fortinata 244 Tab. 2
I. gallaria 244 Tab. 2
I. nelsonaria 244 Tab. 2
I. sp. 244 Tab. 2
I. sp. nov. 244 Tab. 2
I. variabilis 244 Tab. 2
- islands: evolutionary genetics of Asian mammals 69–71 & Fig. 1, 76, 81
- isocitrate dehydrogenase (*Icd-1*, *Icd-2*, E. C. 1.1.1.37) 72
Icd-1
Apodemus 87 App. 2
Niviventer 88 App. 2
Icd-1, *Icd-2* *Microtus* 89 App. 2
- *isoenzymes: *Culex pipiens* population structure and genetic invasion 329
- Isopoda
Asellus aquaticus: *Notonecta* prey 104–105 & Tab. 1
 polymorphism of *Idotea baltica* 45–68 & Figs 1–6 & Tabs 1–5
- Ithomiinae: butterfly behaviour and mimicry 159–160, 162, 165, 172, 174, 177
see also *Melinaea*, *M. ludovicae* *idae*, *M. scylax*, *Tithorea tarricina* 'transparent' mimicry rings
- jacamars (*Galbula* spp.): *Heliconius* predation 174
G. ruficauda 165
- jacanas: sexual selection acting on females 256 Tab. 1, jackknife tests: resolution of kangaroo phylogeny using DNA hybridization 309, 312–313, 318–325 & Figs 2–3
- Jalmenus evagoras*: lycaenid butterfly 210
- Java: *Niviventer* rats 81
- jewelweeds: fit of flowers around the bodies of bees 357–358, 368, 370–372, 377–378
see also *Impatiens capensis* and *I. pallida* 357
- Juanullos* (Solanaceae): *Melinaea* egg-laying host 172
- Juncaceae: insect food-plants 114 Fig. 1, 116, 125 App. 1, 127 App. 2
- Juncaginaceae: insect food-plants 114 Fig. 1, 125 App. 1, 127 App. 2
- *kangaroos: resolution of kangaroo phylogeny using DNA hybridization, 309–328 & Figs 1–3 & Tabs 1–4
 eastern grey *Macropus giganteus* 309, 318–319 & Tabs 2–4, 325
 red (*Macropus rufus* 309, 314 Tab. 1, 317–319 & Tabs 2–4
 'true' 309–311, 322, 324
see also, rat-kangaroos, tree kangaroos and wallabies
- Kappa statistic, Cohen's: cranial variation in mink 296, 302
- Kermisidae (coccids): phytophagous insects 111
- Kitson Mink Farm, Surrey 299 Fig. 2 & Tab. 4
- Kolmogorov-Smirnov tests
 ornament growth, variation and dimorphism in *Vidua* 132
 patterns in brood parasitism 263
 kookaburras (*Dacelo novaguineae*): lizard predators 287
- Korea: *Microtus fortis* 72, 78
- Kruskal-Wallis tests
 genetics of predation 22–23 Tabs 1–2
Heliconius roosting heights 170
- L-leucyl-alanine (*La-1*, E.C. 3.4.11.1) 72
Apodemus 87 App. 2
Microtus 89 App. 2
Niviventer 88 App. 2
- L-leucylglycylglycine (*Lgg*, E.C. 3.4.11.4) 72
- L-phenylalanyl-L-proline (*Pap*, E.C. 3.4.11.5) 72
Apodemus 87 App. 2
- Labiateae: insect food-plants 112 & Tab. 1, 114 Fig. 1, 117, 125 App. 1
- laboratory studies: isopod *Idotea baltica* 46, 50, 52–53 & Fig. 1, 62–64
- lactase dehydrogenase (*Ldh-1*, *Ldh-2*, E. C. 1.1.1.27) 72
Ldh-1: *Apodemus* 87 App. 2
- *ladybirds (Coccinellidae): genetics of predation 17–18
 melanic (*Adalia bipunctata*) 17, 19–20 & Fig. 1, 24 Tab. 2, 26
 seven-spot (*Coccinella septempunctata*: toxic) 19–21 & Fig. 1, 23, 25–26 & Fig. 2
 two-spot: red (*A. bipunctata*) 17–20 & Fig. 1, 22–23 & Tab. 1, 26
- Lagorchestes* (hare wallabies): resolution of kangaroo phylogeny using DNA hybridization 310, 326
- Lagostrophus*: resolution of kangaroo phylogeny using DNA hybridization 326
Lagostrophus fasciatus (banded hare-wallabies) 310
- Lampropholis*: colour patterns in lizards
L. delicata 273–290 & Figs 1–4 & Tabs 1–2
L. guichenoti 277, 286
- *land snails: Madagascan *Tropidophora* 183, 200–202
- Lantana*: orange heliconiine host-plants 166
- Lap*: Madagascan *Tropidophora* (allozyme data) 186, 189 Fig. 2, 205–206 App.
- Larentiinae: patterns in species description 227–228 & Figs 1–2, 232–233 & Tabs 3–4
- Larix*: *Taeparia* host 250 App.
- larvae
Euphilotes enoptes: unpredictable variation in a facultative mutualism 209–223 & Figs 1–4 & Tabs 1–5
Pararge aegeria (speckled wood butterflies) 143, 145–154 & Figs 2–7
- Lasius* spp. tending *Euphilotes enoptes* larvae 213 Tab. 1 latitudes
 brood parasitism patterns 261–262, 264 Fig. 1, 270–272 App.
 colour patterns in a lizard 275–276, 286, 288–289
 plasticity in *Pararge aegeria* 143–145, 147
 leaf-chewing guilds: phytophagous insects 110, 122
 learning, avoidance: genetics of predation 19–23, 26
 least squares linear regression: colour pattern polymorphism in a lizard 276
- leg lengths: sexual dimorphism in *Metellina segmentata* 345–349 & Figs 1–2 & Tab. 1, 351–352
- Lemnaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2

- *Lepidoptera: behaviour and mimicry 159–180 & Figs 1–4 & Tabs 1–4
- *Lepidoptera: lithinine moths on ferns 239–241 & Tab. 1, 247, 250 App.
- *Lepidoptera: *Pararge aegeria* (speckled wood butterflies) 143 157 & Figs 1–8
- Lepidoptera
 - Geometridae: patterns in species description 225–226, 235
 - Impatiens* pollination 378
 - phytophagous insects 111, 122
 - see also, Choreutidae, Chrysomelidae and Curculionidae, micro- and macrolepidoptera
 - unpredictable variation in a facultative mutualism 209–210
 - see also butterflies, macro-, microlepidoptera and moths
- Leptothorax* species: tending *Euphilotes enoptes* larvae 213
 - L. nevadensis* complex 213–214 & Tab. 1
 - L. rugatulus* 213–214 & Tab. 1
- Lever, C. & Poyser, A. D. *Naturalised Animals: the Ecology of Successfully Introduced Species*, (review: D. W. Yalden) 91
- Lichenes: insect food-plants 111, 113–118 & Figs 1–3, 126–127 Apps 1–2
- life forms: insect food-plants 109–127 & Figs 1–8 & Tab. 1
- life history: *Pararge aegeria* (speckled wood butterflies) 143–144, 151
- life-cycle regulation: *Pararge aegeria* (speckled wood butterflies) 143 144, 147–151 & Fig. 2
- Liliaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 120–121, 126–127 Apps 1–2
- Liliales 120–121
- Linaceae: insect food-plants 112–114 & Tab. 1 & Fig. 1, 126–127 Apps 1–2
- Linales 113
- LINKDOS: *Culex pipiens* D statistics 335
- Lithina*
 - = *Petrophora*: evolutionary perspective 246
 - citation as *Tacparia* 250 App.
- *Lithinini (moths): pteridophagy 239, 241–250 & Figs 1–2 & Tabs 1–2 & App
- Littorina*: shell variation 202
 - L. saxatilis* 202
- *Littorinoidea: Madagascan *Tropidophora* 183–184, 195–196
- Livingstonville, Schoharie County: fit of flowers around the bodies of bees 361–362 & Tabs 1–2, 368–369 & Fig. 6, 371, 373–376 Tabs 5–8
- lizards: colour patterns 273 286, 288–289
 - see also *Lampropholis* and Australian scincids
- Lobelia*: pollination
 - L. cardinalis* 379
 - L. siphilitica* 379
- loci
 - Culex pipiens* population structure and genetic invasion 329, 331–333 & Tab. 1, 335–338 & Tabs 3–4
 - see also *Aat*, *Aco*, *Fum*, *Gpd*, *Gpi*, *Idh-1*, *Idh-2*, *Mdh-1*, *Mdh-2*, *Me*, *Odh*, *Pgd*, and *Pgm*
- evolutionary genetics of Asian mammals 72–76 & Tabs 1–2, 78, 87–89 App. 2
- Madagascan *Tropidophora* 186, 200
- log-linear analysis: colour patterns in a lizard 277–278, 283
- log-likelihood ratio 278
- London Natural History Museum
 - geometrid collection 226, 234
 - lithinine moths on fern data 241 Tab. 1
- looper moths, ennomine: pteridophagy 241, 244, 247
- Loranthaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2
- 'lower' plants: insect food-plants 111, 117–118 & Fig. 3
 - see also algae, fungi, lichens and mosses
- LUM (Lumio, Corsica): *Culex pipiens* sample site 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2
- Lutra lutra* (otters): dimorphic comparisons 294
- Lycaenidae
 - Hypochrysops theon*: fern-feeding 240
 - unpredictable variation in a facultative mutualism 209 211, 219 221
 - Lycaena tityrus* 219
 - lycaenid-ant interactions 210, 214
 - see also *Euphilotes enoptes*, *Icaricia acmon*, and *Jalmenus evagoras*
- Lythraceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2
- macrolepidoptera: phytophagous insects 110, 122
- macro-moths 117
- *Macropodidae: resolution of kangaroo phylogeny using DNA hybridization 309–328 & Figs 1–3 & Tabs 1–4
- Macropodinae 309–319 & Tabs 1–4 & Fig. 1, 323–326
 - see also *Dorcopsulus*, *Macropus* and *Setonix*
- Macropus*: resolution of kangaroo phylogeny using DNA hybridization 309 311, 313, 315, 318–319, 324–326
 - M. bernardus* 325
 - M. giganteus* (eastern grey kangaroos) 309, 318–319 & Tabs 3–4, 325
 - M. parryi* (Parry's wallabies) 309, 312, 314 Tab. 1, 317–321 & Tabs 2–4 & Fig. 2, 325
 - M. robustus* (wallaroos) 309, 318–319 & Tabs 3–4, 325
 - M. rufogriseus* (red-necked wallabies) 309, 314–315 Tab. 1 & Fig. 1, 317–321 & Tabs 2–4 & Fig. 2
 - M. rufus* - *M. robustus* 325
 - M. rufus* (red kangaroos) 309, 314 Tab. 1, 317–319 & Tabs 2–4, 320 321 Fig. 2, 325
 - Macropus* - *Wallabia* grouping 319
- Madagascar
 - biogeographic region (5) 226, 228–231 & Tabs 1–2 & Fig. 3, 234 & Fig. 8
 - Tropidophora* (snails) 183 208 & Figs 1–8 & Tabs 1–2 & App
- Maddocks, Linda, Bone, Quentin and Raynor, Jeremy M. V., editors *Mechanics and Physiology of Animal Swimming*, (review: H. D. Jones) 385
- Madeira: *Pararge aegeria* (speckled wood butterflies) 143–143, 145–146, 153 154 & Fig. 7
- Magnoliidae: insect food-plants 118–119 & Fig. 4
- Magnoliopsida (Dicotyledonae): larval hosts for Lithinini 250 App.

- magpies, Australian (*Gymnorhina tibicen*): lizard predators 287
- Mahalanobis distances: cranial variation in mink 297–302 & Tabs 3–4 & 6–7
- malate dehydrogenase (*Mdh-1*, *Mdh-2*, E.C. 1.1.1.37.) 72
- male phase: fit of flowers around the bodies of bees 358–360 & Fig. 1, 367–368
- males
- African whydahs (*Vidua*) 129–140 & Tabs 1–7
 - colour patterns in a lizard 273–275, 277–289 & Figs 1–2 & 4–5 & Tab. 1
 - cranial variation in mink 294–295 & Tab. 1, 298, 300–305 & Figs 3–4 & Tabs 6 & 9
 - Heliconius*: mating behaviour 166–167, 176
 - insect prey experiments: Japanese quail 19–20, 23
 - Madagascan *Tropidophora* 188, 190–201 & Figs 3–8 & Tab. 2
 - Pararge aegeria* (speckled wood butterflies) 146–147 & Fig. 1, 149 Figs 3–4, 150, 152, 154 Figs 6–7
 - polymorphism of the isopod *Idotea baltica* 45, 47–65 & Tabs 1 & 3 & 5 & Figs 1 & 3–6
 - neomales 48–49, 65
 - sexual dimorphism in *Metellina segmentata* 345–352 & Tab. 1 & Fig. 2
 - sexual selection processes 251–252, 254–258 & Fig. 2
- malic enzyme (*Me-1*, *Me-2*, E.C. 1.1.1.40.) 72
- Me-1*, *Me-2*
 - Microtus* 89 App. 2
 - Niviventer* 88 App. 2
 - Me-2*: *Apodemus* 87 App. 2
- Malvaceae: insect food-plants 112–114 & Tab. 1 & Fig. 1, 119, 126–127 Apps 1–2
- Malvales 113, 119
- mammals
- cranial variation in mink 293–294
 - sex-linked colour polymorphism 274
 - small: evolutionary genetics
 - Asia 69–89 & Figs 1–3 & Tabs 1–4 & Apps 1–2
 - North America 70
- Manly's index (B): variation in predator preference 96
- Mann-Whitney values
- cranial variation in mink 297
 - genetics of predation 22–24 & Tabs 1–3
- MANOVA (multivariate analysis of variance): colour patterns in a lizard 276, 281 & Tab. 1
- mark-recapture program: butterfly behaviour and mimicry in *Heliconius* 163, 167
- Markea* (Solanaceae): *Melinaea* egg-laying host 172
- Markov chain method: probability computations 333–335
- *marsupial evolution: resolution of kangaroo phylogeny using DNA hybridization 309–310, 323–324
- marsupials (Argyrolagidae): Pleistocene 41
- *mate choice: sexual selection acting on females 251–252, 254, 258
- mate-searching activity: polymorphism of the isopod *Idotea baltica* 45, 47–48
- mate choice experiment 46, 51–52, 64
- material and methods
- butterfly behaviour and mimicry in *Heliconius* 162–163
 - colour patterns in a lizard 275–278
 - convergence in harvester ant assemblages 31–35 & Fig. 1 & Tab. 1, 44 App.
 - cranial variation in mink 295–296 & Tab. 1
 - Culex pipiens* population structure and genetic invasion 331–335 & Tab. 1 & Fig. 1
 - evolutionary genetics of Asian mammals 69, 71–73
 - Madagascan *Tropidophora* 185–188 & Fig. 1 & Tab. 1
 - plasticity in *Pararge aegeria* 143, 145–146
 - polymorphism of the isopod *Idotea baltica* 45–46, 48–52 & Tab. 1
 - relationships between polyaxonid and monaxonid spicules in Demospongiae 1–3
 - resolution of kangaroo phylogeny using DNA hybridization 311–313
 - sexual dimorphism in *Metellina segmentata* 346–348 & Fig. 1 & Tab. 1
 - unpredictable variation in a facultative mutualism 211–213
 - see also methods
- materials: fit of flowers around the bodies of bees 357–359 & Fig. 1
- mating success (MS): sexual selection acting on females 251–252, 255–258 & Fig. 2 & Tab. 1
- MDH (malate dehydrogenase)
- Culex pipiens* population structure and genetic invasion 331
 - Mdh* allele 336
 - Mdh-1* locus 332–334 & Tab. 1, 334, 336–338 & Tabs 3–4
 - Mdh-2* locus 332–334 & Tab. 1, 337–338 & Tabs 3–4
 - Madagascan *Tropidophora* (allozyme data): *Mdh* 186, 189 ff. 2, 205–206 App.
- ME (malic enzyme): *Culex pipiens* population structure and genetic invasion 331
- Me* locus 333–334 & Tab. 1, 337–338 & Tabs 3–4
- mealworms: quail prey experiments 20–24 & Fig. 1 & Tabs 2–3, 26
- dark brown lesser (*Alphitobius diaperinus*): controls (C) 19–20 & Fig. 1, 26
 - light brown (*Tenebrio molitor*) 19
- means
- convergence in harvester ant assemblages 37–38 & Figs 2–4 & Tab. 3
 - fit of flowers around the bodies of bees 361–363 & Tab. 2, 365–367 Figs 3–5, 372–373 & Tab. 5, 376 Tab. 8
- measurements
- errors (MEs), cranial variation in mink 296, 303
 - fit of flowers around bees 363
 - insect prey experiments
 - fear 17, 19, 21–22
 - feeding responses 21
 - prey preferences 95–96
- Mechanics and Physiology of Animal Swimming*, edited by Linda Maddocks, Quentin Bone and Jeremy M. V. Raynor. (review: H. D. Jones) 385
- *Mediterranean ecology: patterns in brood parasitism 261–267 & Tab. 1
- Mediterranean, western: population structure and genetic invasion 350
- megaphanerophytes: insect food-plants 111
- Melbourne (M): *Lampropholis delicata* 276, 279 Fig. 1
- Meles meles* (badgers): dimorphic comparisons 294

- Melinaea*: butterfly behaviour and mimicry 160
M. ludovica idae 165
M. scylax 164–165 & Fig. 1, 172
- Melophorus*: Australian harvester ants 40
- membracid-ant association 210
- Menyanthaceae: insect food-plants 114 Fig. 1, 119, 126–127 Apps 1–2
- Meranoplus*: Australian harvester ants 40
- Mesodon elevatus* and *M. zaletus* (snails): Tenesse, U.S.A. 200
- mesophanerophytes: insect food-plants 111
- Mesophlebia* (*Entomopteryx*): *Lithinini* larval hosts 250 App.
- Messor*: North American harvester ants 40
- meta-analysis: fit of flowers around the bodies of bees 374 Tab. 6
- Metellina segmentata*: sexual dimorphism 345–354 & Figs 1–3 & Tab. 1
- methods
case study of variation in predator preference 104
fit of flowers around the bodies of bees 359–361 & Tab. 1
genetics of predation 17–22 & Fig. 1
insect food-plants, life form and taxonomy 109–112
ornament growth variation, and dimorphism in *Vidua* 129, 131–132
patterns in brood parasitism 262–263
see also material and methods
- Metidae: sexual dimorphism in *Metellina segmentata* 345
- '*Metrocampa*' (*Pteridium*): *Lithinini* larval hosts 250 App.
- Mexico
harvester ant assemblages 32 Fig. 1
Heliconius charitonia 175
Neotropics 226
- MGPCAs see multiple group principal components analyses
- mice
see *Apodemus*
laboratory: analogous to ranch mink 305
Mus: mandibular genetics 294–295
- Microciona gradalis*: spicules in Demospongiae 3, 6
- microhabitats: colour patterns in a lizard 273, 285, 287, 289
- Microlepidoptera: phytophagous insects 115
- microphanerophytes: insect food-plants 110
- Microtus* (voles): evolutionary genetics of Asian mammals 72, 75–76 & Tab. 2, 78–79 & Fig. 2
M. californicus: United States 72, 75–76 & Tab. 2, 78–79 & Fig. 2, 86 App. 1, 89 App. 2
M. fortis: Korea 72, 75–76 & Tab. 2, 78–79 & Fig. 2, 86 App. 1, 89 App. 2
M. kikuchii 69–79 & Tabs 1–4 & Fig. 2, 81–83, 86 App. 1, 89 App. 2
- midges, wheat blossom (*Sitodiplosis mosellana*): staggered diapause 122–123
- *migration: *Culex pipiens* population structure and genetic invasion 329–331, 335, 338–341 & Tab. 5
- mimicry rings
Heliconius butterflies 159–180 & Figs 1–4 & Tabs 1–4
'transparent': ithomiine butterflies 162, 165
see also
Batesian and M/V811lerian mimicry
'blue', 'red', 'orange', 'tiger' and 'yellow' rings
- mink, American (*Mustela vison*): cranial variations 293–307 & Figs 1–4 & Tabs 1–9
- mint, Iberian: variations in pollinators 210
- Miocene
Andes Mountain Range 39
fossil records: Macropodinae 310, 323–324
- mites: phytophagous arthropods 109–113 & Tab. 1, 121, 123
- models
butterfly behaviour and mimicry 159–161, 176–177
see also *Melinaea*
chemical facilitation: lithinine moths on ferns 239–240, 242
colour patterns in a lizard 274, 278–279
fit of flowers around the bodies of bees 375 Tab. 7 n.
patterns in species description 235–236
sexual dimorphism in *Metellina segmentata* 345, 349
sexual selection acting on females 252, 255–256 & Fig. 2, 258
variation in predator preference 93–95 & Fig. 1
bimodal 96–103 & Figs 2–4
normal distribution 96, 101–103 & Figs 5–6
- *molecular phylogeny: resolution of kangaroo phylogeny using DNA hybridization 310
- monactines: relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 11–12
- Monarda*: pollination of trimmed flowers 380
- monaxonid spicules: relationships between them and polyaxonid spicules in Demospongiae 1–2, 7, 11–12
- monocotyledons: insect food-plants 109, 113–114, 117, 119–122 & Fig. 8
- Monomorium*: Australian harvester ants 40
- monomorphism: sexual selection acting on females 251, 258
- Monte desert, Argentina: harvester ants 39–41
- Moorea, Society archipelago: dispersal of resistance genes in *Culex pipiens* 339
- Morisita's index (I_m): assessment larval aggregation 213, 216, 218 Tab. 3
- *morphological variation: Madagascan *Tropidophora* 183–184
- *morphological variation: whydahs (*Vidua*) 129–141 & Tabs 1–7
- morphology
cranial variation in mink 294
fit of flowers around the bodies of bees 355–373 & Figs 1 & 4–8 & Tabs 2–4, 375–380 & Tabs 7–8
relationships between polyaxonid and monaxonid spicules in Demospongiae 1–2, 7, 10, 12
see also acanthostyles, actines, canals, desmas, desmoids, oxeas, sphaeroclons, spicules, spines, tubercles and tylostyles
- morphs, colour
isopod *Idotea baltica* 45–58 & Figs 1–3 & Tabs 2–5, 61–65
determination 49
fine-mottled: *maculata* 48–49, 52–57 & Figs 1–3 & Tabs 2–3, 62–63, 65
striped, *bilineata* and *bilineata-lineata* 49
two-striped: *bilineata* 48
uniform (light and/or dark): *uniformis* 45, 48–64 & Figs 1–3 & 5–6 & Tabs 2–5

- white-spotted (light and/or dark): *albafusca* 45, 47–62 & Figs 1–3 & 5–6 & Tabs 2–4, 64
- mortality: differential predation on sexes of isopod *Idotea baltica* 45, 50–51 & Tab. 1, 63–64
- *mosquitoes (*Culex pipiens*): population structure and genetic invasion 329–343 & Figs 1–2 & Tabs 1–5
- moths
- arctiine (Pericopinae): tiger patterns 172
 - flower fits 356
 - hawkmoth pollination: *Viola cazorlensis* 379
- Geometridae: patterns in species description 225–237 & Figs 1–8 & Tabs 1–4
- lithinine: pteridophagy 239–250 & Figs 1–2 & Tabs 1–2 & App.
- see also looper moths
- moult: ornament growth, variation and dimorphism in *Vidua* 129, 131, 139
- mountaintops: evolutionary genetics of Asian mammals 69–70, 72–73, 77, 82–83
- see also Mt Guanshan and Mt Yushan
- Mpi*: Madagascan *Tropidophora* (allozyme data) 186, 189 Fig. 2, 205–206 App.
- Mt Guanshan: Taiwanese small mammals 72, 78, 82, 85–86 App. 1
- Mt Yushan: Taiwanese small mammals 72, 78, 82
- mtDNA geneology: comparison of *Heliconius erato*- and *H-melpomene* groups 177
- Müllerian mimicry
- butterfly behaviour 159–161, 176–178 & Fig. 4
 - variation in predator preference 106
- multiple group principal component analyses (MGPCAs)
- cranial variation in mink 296, 298
 - sexual dimorphism in *Metellina segmentata* 347–349 & Figs 2–3
- multivariate analyses
- insect food-plants 110
 - sexual dimorphism in *Metellina segmentata* 345–352
 - see also multiple group principal component analyses (MGPCAs)
- *multivariate morphometrics: cranial variation in mink 293–294
- Murdoch's index *c*: variation in predator preference 96, 98
- Mus*: mandibular genetics 294–295
- Musci (mosses): insect food-plants 111, 113–118 & Figs 1–3, 126–127 Apps 1–2
- Musotiminae (lepidopteran subfamily): fern associations 241 Tab. 1
- Mustela*
- M. putorius furo* (ferrets): domestication 305
 - M. putorius* (polecats): domestication 305
 - M. vison* (American mink): cranial variations 293–306 & Figs 1–4 & Tabs 1–9
- Mustelidae: cranial variation in mink 293–295, 305
- see also ferrets, polecats, stoats and weasels
- *mutualism: *Euphilotes enoptes* and ants 209–210, 219–221
- Mycetophilidae: phytophagous insects (fungi) 110
- Myricaceae
- insect food-plants 114 Fig. 1, 117–118, 126–127 Apps 1–2
 - larval hosts for *Tacparia* 250 App.
 - Myrica* 247, 250 App.
 - see also *Comptonia*
- Myrmicinae
- harvester ants 40
 - tending *Euphilotes enoptes* larvae 213 Tab. 1
 - Myrmica* nr. *lobifrons* 213 Tab. 1
 - see also *Leptothorax*
- Mytilus edulis*: epibiont background for isopod *Idotea baltica* 48
- Myxilla rosacea*: spicules in Demospongiae 3, 6
- Nadagara* (loopers moths): fern associations 241, 245 Fig. 2
- Namibia: ornament growth, variation and dimorphism in *Vidua* 131, 134–135 Tabs 2–5
- nanophanerophytes: insect food-plants 111
- Naturalised Animals: the Ecology of Successfully Introduced Species*, by C. Lever & A. D. Poyser (review: D. W. Yalden) 91
- Nearctic: biogeographic region (1) 226, 228–231 & Tabs 1–2 & Fig. 3, 234 Fig. 8
- 'nearest neighbours': insect food-plants 109, 112, 123, 125–126 App. 1
- Nectarinia johnstoni* (sunbirds): sexual signalling 139
- Nei's genetic distances
- evolutionary genetics of Asian mammals 72, 75–76 & Tabs 2–3, 82
 - Madagascan *Tropidophora* 186, 189 Fig. 2, 199–200
- Nelson consensus tree: Madagascan *Tropidophora* 186, 189 & Fig. 2
- nematode *Skjabinjylus nasicola* infection: American mink 303
- neophobia: genetics of predation 18, 20
- Neotropics
- biogeographic region (2) 226, 228–231 & Tabs 1–2 & Fig. 3, 234–235 & Fig. 8
 - see also Mexico and South America
 - patterns in brood parasitism 270 App.
- Netherlands
- Japanese quail 19
 - patterns in brood parasitism 263–266 & Figs 2–3
 - see also Nunspeet
- New Guinea: resolution of kangaroo phylogeny using DNA hybridization 310
- forest wallaby (*Dorcopsulus vanheurni*) 309, 311, 313, 323–324
- New York: fit of flowers around the bodies of bees 361 & Tab. 1
- see also Albany, Freehold, Livingstonville, Long Island, Rensselaerville and Trout Creek
- New Zealand
- biogeographic region (9) 227–231 & Tabs 1–2 & Fig. 3, 234–235 & Fig. 8
 - fossils: relationships between polyaxonid and monaxonid spicules in Demospongiae 3, 7–8 & Fig. 4
 - moths: *Ischalis* pteridophyte associations 244–246 & Tab. 2
 - see also *Ischalis*, *Sarisa* and *Sestra*
- Niniventer*: evolutionary genetics of Asian mammals 69–71, 74–76 & Tab. 2, 78–81 & Figs 2–3
- N. bukit*: China 72, 74–76 & Tab. 2, 78–80 & Figs 2–3, 86 App. 1, 88 App. 2
 - N. coxingi* 72, 74–76 & Tabs 2–3, 78–80 & Figs 2–3, 86 App. 1, 88 App. 2
 - N. culturatus* 69–77 & Tabs 1–2 & 4, 78–82 & Figs 2–3, 85 App. 1, 88 App. 2

- Noctuidae
 lepidopteran family: fern associations 240–241 & Tab. 1
 Noctuoidea (superfamily) 241 Tab. 1
 species-rich Lepidoptera 226
- North America
 eastern: *Impatiens capensis* and *I. pallida* 357
 harvester ants 29–33 & Fig. & Tab. 1, 35–41 & Tabs 2–4 & Figs 2–3, 69–77 & Tabs 1–2 & 4, 78–82 & Figs 2–3, 85 App. 1, 88 App. 2
Mustela vison (American mink) 294
 Nearctic biogeographic region (1) 226
Peromyscus maniculatus: adaptation to hypoxia 82
see also Canada and United States (USA)
- Norway: American mink 294–296 & Tab. 1, 300–301 & Fig. 4 & Tab. 7
- **Notonecta*: variation in predator preference 93, 104–106 & Tab. 6
N. glauca 104
- Novomessor*: North American harvester ant assemblages 40
- Nunspeet, Netherlands: patterns in brood parasitism 263–266 & Figs 2–3
- Nymphaeaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2
 Nymphales 119
- Nymphalidae
Pararge aegeria (speckled wood butterflies) 143
 peritridophagy (erroneous record) 240
- oaks: insect food-plants 122
- Ochroma*: Corcovado secondary forest 163
- ODH (octanol dehydrogenase): *Culex pipiens* population structure and genetic invasion 331
Odh locus 333 Tab. 1, 335, 337–338 Tabs 3–4
- Oecophoridae (lepidopteran family): fern associations 241 Tab. 1
- Oenochrominae: patterns in species description 227–229 & Figs 1–2 & Tab. 1, 232 = 233 & lts 3–4
- offspring: sexual selection acting on females 251–252, 254–255, 257
- Ohta indices: population structure and genetic invasion 335–336
- OLB (Olbia, Sardinia): *Culex pipiens* sample site 332 334 & Tab. 1 & Fig. 1, 336 Tab. 2, 338
- Oleaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 119, 126–127 Apps 1–2
- Onagraceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 126–127 Apps 1–2
- ontogeny: polyaxonid and monaxonid spicules in Demospongiae 1–2, 5–6, 11–12
- Onychogalea* (nail-tailed wallabies): resolution of kangaroo phylogeny using DNA hybridization 310, 326
- operational sex ratio (OSR): sexual selection acting on females 252, 258
- Ophraella* (chrysomelid beetles): insect-plant interactions 248
- opisthosomata: sexual dimorphism in *Metellina segmentata* 347–349 & Figs 1–2 & Tab. 1, 351
- 'orange' mimicry rings: butterflies 162, 165–166, 176
see also *Agraulis vanillae*, *Dione juno*, *Dryadula phaetusa*, *Dryas iulia*, *Eueides aliphera*, *E. lybia* and *E. vibilia*
- orb-weaving spiders: sexual dimorphism in *Metellina segmentata* 345–346, 351–352
- Orchidaceae
 insect food-plants 114 Fig. 1, 126–127 Apps 1–2
 Orchidales 121
 pollination success 378
- orders: insect food-plants 109–110, 119, 122–123
- organophosphates (OPs): *Culex pipiens* population structure and genetic invasion 329–330, 340–341
- Orhisaari: isopod *Idotea baltica* 49–50, 52 57 & Figs 1 3 & Tabs 2 3, 62
- Orient
 component: evolutionary genetics of Asian mammals 80 81
 region: patterns in brood parasitism 270 271 App.
- *ornament size: whydahs (*Vidua*) 129–131, 133–140 & Tabs 4–7
- *ornaments: sexual selection acting on females 251, 253–254 & Fig. 1, 257–258
- Orobanchaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2
- Osphranter*: resolution of kangaroo phylogeny using DNA hybridization 325
- otters (*Lutra lutra*): dimorphic comparisons 294
- outgroups: resolution of kangaroo phylogeny using DNA hybridization 309, 311, 313–323 & Tabs 1–2 & Fig. 2
- Oxalidaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2
- oxeas: relationships between polyaxonid and monaxonid spicules in Demospongiae 2, 6, 9 Fig. 6
- Pachymelania*: shell variation 202
- Pachyrhabda* (Lepidoptera): fern associations 241 Tab. 1
- pademelons (*Thylogale*): resolution of kangaroo phylogeny using DNA hybridization 309–311, 315
- Paesia*: moth associations 242, 245–246 & Fig. 2, 250 App.
- pairwise comparisons
 continents: convergence in harvester ant assemblages 35–36 & Tab. 2
 evolutionary genetics of Asian mammals 75–76 & Tab. 2
- Palaeartic
 component: evolutionary genetics of Asian mammals 70, 80 81
 patterns in brood parasitism 262–264 & Fig. 1, 271 App.
see also Eastern and Western
- palates: cranial variation in mink 293–295, 302
- palatability: genetics of predation 18–21, 26
- Panama: arctine moths (tiger patterns) 172
- Panotima* (Lepidoptera): fern associations 241 Tab. 1
- Papaveraceae: insect food-plants 113–114 & Fig. 1, 117, 119, 126 127 Apps 1 2
- Papilionaceae: insect food-plants 112 & Tab. 1, 114 Fig. 1, 117, 126–127 Apps 1–2
- Papilionidae: host-plant relationships 247–248
- Papilionoidea 240
- **Pararge aegeria* (speckled wood butterflies) 143 155 & Figs 1–7
P. a. aegeria 143, 146
P. a. tircis 143, 145, 147–151 & Figs 2–5

- Pararge xiphia* (speckled wood butterflies): Madeiran endemic 145
- parasitism: patterns in brood parasitism 261–272 & Figs 1–2 & Tab. 1 & App.
- parental care: sexual selection acting on females 251–252, 255–258 & Tab. 1
- parental expenditure (PE) 256 Fig. 2
- passerine birds: sexual selection acting on females 258
- Passiflora*: *Heliconius* larval food plants 167
- P. ambigua* 167
- P. piltieri* 166
- P. spp.* 161–162
- Passifloraceae 161–162
- 'pastel' (PA): ranch mink pelts 297–299 & Figs 1–2 & Tabs 3–4, 304
- Patagonian Andes: Lithinini 241
- patterns
- brood parasitism in bees 261–262, 265, 267
 - diversification and genetic population structure of small mammals in Taiwan 69–72, 78, 80–81
 - isopod *Idotea baltica* backgrounds 46–48, 62–63
 - Lampropholis delicata* polymorphisms 273–276, 279–280, 284–289
 - lithinine moths on ferns 239–244 & Tab. 1, 246–248
 - sexual selection acting on females 251–252, 255–256 & Tab. 1, 258
 - species descriptions 225–237 & Figs 1–8 & Tabs 1–4
 - see also colour and latitudes
- PAUP
- evolutionary genetics of Asian mammals 72–73, 78
 - lithinine moths on ferns 246
- Peasia* (Dennstaedtiaceae): moth associations 242, 245 Fig. 2
- *pelt colours: cranial variation in mink 293–295, 297–299 & Tabs 3–4 & Fig. 2, 303–305
- see also 'pastel', 'sapphire' and 'silverblu'
- penes: Madagascan *Tropidophora* 183–184, 186, 188, 195–201 & Figs 6–8
- Pep-G*: Madagascan *Tropidophora* (allozyme data) 186, 189 Fig. 2, 207–208 App.
- Pep-TI*: Madagascan *Tropidophora* (allozyme data) 186, 189 Fig. 2, 207–208 App.
- Pep-T2*: Madagascan *Tropidophora* (allozyme data) 186, 189 Fig. 2, 207–208 App.
- Perca fluviatilis* (perch): common predator of isopod *Idotea baltica* 45, 47, 50–51, 57–59 & Tab. 5 & Fig. 4, 62, 64
- Peregrine Fur Farm, Devon, Britain: mink 294–295, 297–299 & Tabs 2–4, 304
- Pericopinae (arctiine moths): tiger patterns 172
- Peromyscus*
- P. californicus*: gene flow 82
 - P. maniculatus*: adaptation to hypoxia 82
- Peru: butterfly behaviour and mimicry 161, 165, 174
- Petrogale* (rock wallabies): resolution of kangaroo phylogeny using DNA hybridization 309–311, 318–319, 324, 326
- P. mareeba* 315
- P. sharmani* 314–315 & Tab. 1, 317 Tab. 2, 320–321 Fig. 2
- P. xanthopus* 313–315 & Tab. 1 & Fig. 1, 317 Tab. 2, 320–321 Fig. 2
- Petrophora* (*Dryopteris*, *Pteridium* and *Thelypteris*): *Ischalis* host associations 250 App.
- PGD (phosphoglucosmutase)
- Culex pipiens* population structure and genetic invasion 331
 - Pgd* locus 333 Tab. 1, 337–338 Tabs 3–4
 - Madagascan *Tropidophora* (allozyme data): *6Pgd* 186, 189 Fig. 2, 207–208 App.
- PGM (phosphoglucosmutase)
- Culex pipiens* population structure and genetic invasion 331
 - Pgm* locus 333 Tab. 1, 336–338 & Tabs 3–4
 - Madagascan *Tropidophora* (allozyme data): *Pgm* 186, 189 Fig. 2, 207–208 App.
- Phalangeridae: resolution of kangaroo phylogeny using DNA hybridization 309, 323
- see also *Trichosurus*
- phalangers, brush-tailed (*Trichosurus vulpecula*): resolution of kangaroo phylogeny using DNA hybridization 309, 313
- phalaropes: sexual selection acting on females 256 Tab. 1,
- phanerophytes (trees and shrubs): insect food-plants 111, 114–116 & Fig. 2
- Pheidole*: Australian, North and South American harvester ant assemblages 39–40
- P. sp.* (South America) 44 App.
- phenology: plasticity in *Pararge aegeria* 143–146, 151
- Philadelphia: Academy of Natural Sciences (ANSP): Madagascan *Tropidophora* 201, 205–208 App
- Philedia* (*Pteridium*): *Ischalis* host association 250 App.
- Philippines
- faunistic affinities 81
 - Haplomycteris* bats 83
- Phorbas fictitus*: spicules in Demospongiae 3, 6
- phosphoglucosmutase (*Pgm*, E.C. 2.7.5.1.) 72
- Apodemus* 87 App. 2
- Microtus* 88 App. 2
- Niviventer* 89 App. 2
- 6-phosphoglucosmutase dehydrogenase (*6-Pgd*, E.C. 1.1.1.44) 72
- Apodemus* 87 App. 2
- Microtus* 89 App. 2
- Niviventer* 88 App. 2
- *photoperiod: plasticity in *Pararge aegeria* 143–144, 146–149 & Fig. 1, 151–153
- PHYLIP package: resolution of kangaroo phylogeny using DNA hybridization 312
- Phyllobrotica* (chrysomelid beetles): insect-plant interactions 248
- *phylogeny: convergence in harvester ant assemblages 29–31, 40–41
- phylogeny
- Asian and Taiwanese small mammals 69, 71–72 & Fig. 1, 78, 80–81 & Fig. 3
 - kangaroos: resolution using DNA hybridization 310, 315–319 & Tabs 2–4, 324–326
 - see also trees
 - lithinine moths on ferns 239–248 & Figs 1–2
- 'Phymatodes' (= *Polypodium*): *Sarisa* citation 250 App.
- Phymatosorus* (Polypodiaceae): moth associations 242, 244–246 & Tab. 2 & Fig. 2, 250 App.
- Phytophagous Insects Data Bank (PIDB) 110, 112 Tab. 1, 116, 123
- Picea*: *Thallophaga* host 250 App.

Pinaceae

- insect food-plants 114–115 & Fig. 1, 117, 126–127 Apps 1–2
- larval hosts for Lithiniinae 250 App.
- Pinopsida 247, 250 App.
- see also *Abies*, *Larix*, *Picea*, *Pinus*, *Pseudotsuga* and *Tsuga*; Cupressaceae

Pinus

- Reecer and Tarpiscan Creeks: *P. ponderosa* 211–212
- Thalophaga* host 250 App.
- pipefish: sexual selection acting on females 256 Tab. 1,
- plant families: insect food 109–127 & Figs 1–8 & Tabs 1 & Apps 1–2
- see also 'lower' plants
- Plantaginaceae: insect food-plants 112 & Tab. 1, 114 Fig. 1, 117, 126–127 Apps 1–2
- plasticity, life history: *Pararge aegeria* (speckled wood butterflies) 143 144
- Platanaceae: insect food-plants 114 Fig. 1, 118, 126–127 Apps 1–2
- Pleistocene
 - marsupials: Argyrolagidae 41
 - Taiwan connection with the Asian continent 70, 81
- Pliocene: Taiwan connection with the Asian continent 70
- Ploceidae (*Vidua*): ornament growth variation and dimorphism 129
- Plumbaginaceae: insect food-plants 114–116 & Fig. 1, 119, 126–127 Apps 1–2
- Pneumatopteris*: *Ischalis* host association 244–245 Tab. 2 & Fig. 2, 250 App.
- Poa annua* (host grass): *Pararge aegeria* (speckled wood butterflies) 145–147
- Poecilosclerida: spicules in Demospongiae 1, 11 12
- POG (Santa Maria di Poggio, Corsica): *Culex pipiens* sample site 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2
- Pogonomyrmex*: harvester ant assemblages
 - North America 40
 - South America
 - Argentina 39
 - P. breviparbis* 44 App.
 - P. cunicularius* 44 App.
 - P. inermis* 44 App.
- polecats (*M. putorius*): domestication 305
- Polemoniaceae: insect food-plants 114 Fig. 1, 119, 126–127 Apps 1–2
- Polemonium* floral traits: pollination 379–380
- *pollen deposition: fit of flowers around the bodies of bees 355–361, 363–380 & Tabs 4–8 & Figs 4–8
- *pollen removal: fit of flowers around the bodies of bees 355–361, 363–380 & Tabs 4–8 & Figs 4–8
- pollination: fit of flowers around the bodies of bees 357, 378–380
- pollinators: fit of flowers around the bodies of bees 355–358, 365, 371, 377–380
- *polyaxonid spicules: relationships between them and monaxonid spicules in Demospongiae 1–2, 5–6, 10–12
- Polygalaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2
- Polygonaceae
 - Eriogonum*: *Euphilotes enoptes* host 211

insect food-plants 112 Tab. 1, 114 Fig. 1, 117,

126–127 Apps 1–2

polygyrid snails: Madagascan *Tropidophora* comparisons 200

Mesodon elevatus and *M. zaletus* 200

*polymorphism: colour patterns in a lizard 273–276, 279–281, 286–290

polymorphisms

Culex pipiens population structure and genetic invasion 335 336

differential predation affects colour of isopod *Idotea baltica* 45–47

Polyommatainae (subfamily): ant calls 211

Polyodiaceae

insect food-plants 114 Fig. 1, 116–117, 126–127 Apps 1–2

see also *Pteridium*

moth associations 242, 244–246 & Tab. 2 & Fig. 2, 248, 250 App.

Polypodium diversifolium: citation as *Phymatosaurus* 250 App.

Polystichum (Dryopteridaceae): moth associations 242, 244–245 Tab. 2 & Fig. 2, 250 App.

Pomatiasidae: Madagascan *Tropidophora* 183–184, 195, 202

Pomatias 195–196

populations

American mink 293 295, 303, 305

see also feral and ranch

consequences of variation in predator preferences 93–107 & Figs 1–6 & Tab. 1

differentiation: *Culex pipiens* 334–335, 337–338 & Tab. 3, 340–341

evolutionary genetics of Asian mammals 69–70, 73–83 & Tabs 1–2 & 4 n. & Figs 2–3, 87–89 App. 2

subpopulations 76–77 & Tab. 3, 82–83

fit of flowers around the bodies of bees 355, 358, 362–365, 368–378 & Figs 5–8 & Tabs 5–8, 380

Lampropholis delicata 273 281 & Figs 1–2, 286–290

Madagascan *Tropidophora* 183, 186–195 & Tabs 1–2, 200–201

ornament growth, variation and dimorphism in *Vidua* 129–131, 136, 140

Pararge aegeria (speckled wood butterflies) 143–153 & Fig. 1 & 3 5

Porifera: relationships between polyaxonid and monaxonid spicules in Demospongiae 1–15 & Figs 1–7

Portulacaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2

postorbital constrictions: cranial variation in mink 293, 295, 302, 305

Potamogetonaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1 2

Potoroinae: resolution of kangaroo phylogeny using DNA hybridization 309, 315, 323

see also

Aepyprymnus

rat kangaroos

Potorous tridactylus: resolution of kangaroo phylogeny using DNA hybridization 315

precipitations: convergence in harvester ant assemblages 31, 33–34, 39

- intervals 33 & Tab. 1, 35–37 & Tab. 2 & Figs 2–3, 39
- *predation: butterfly behaviour and mimicry 159–161, 165, 172, 174–175, 177–178 & Fig. 4
- *predation: colour patterns in a lizard 273, 275–277, 286–287, 289
- predation
- polymorphism of the isopod *Idotea baltica* 45–52 & Tab. 1, 55–65 & Tabs 4–5
 - variation in preferences 93–107 & Figs 1–6 & Tab. 1
- *preference, predator: population consequences 93–107 & Figs 1–6 & Tab. 1
- measurement 93–96
 - variation 96–97
- Prenolepis*: South American harvester ant assemblages 40
- P. longicornis* 44 App.
- preserved specimens: *Lampropholis delicata* 273–274, 276–282 & Figs 1–3 & Tabs 1–2, 288–289
- prey colouration: isopod *Idotea baltica* 46–47, 62
- *prey recruitment: genetics of predation 17–21 & Fig. 1
- times (PRT) 18–24 & Tabs 1–2
- Primulaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 126–127 Apps 1–2
- principal components (PCs)
- cranial variation in mink 296–299 & Tabs 2 & 5, 302–303 & Tabs 8–9
 - ornament growth, variation and dimorphism in *Vidua* 136, 138 Tab. 7
 - sexual dimorphism in *Metellina segmentata* 345, 348–350 & Fig. 3
 - PC1 (shape) 345, 348–352 & Figs 2–3
 - PC2 (size) 345, 348–352 & Figs 2–3
- principal coordinates analysis: insect food-plants 109–115 & Figs 1–2, 117–122 & Figs 3–8
- Czelkanowski coefficient 111–112
- pro-apostatic selection: variation in predator preference 93, 96, 99–100, 103, 106
- probabilities, combinatorial: orders of *Heliconius* mimicry rings 169–170, 172
- procedures *see* sampling and statistical
- *Prosobranchia: Madagascar *Tropidophora* 183, 201–202
- see also Cerithium* 202
- prosomata: sexual dimorphism in *Metellina segmentata* 345–350 & Figs 1–2 & Tab. 1, 352
- Pseudomyrmecinae: convergence in harvester ant assemblages 40
- Pseudomyrmex*: South American harvester ants 40
- P. denticollis* 44 App.
- Pseudotsuga*
- Reecer Creek: *P. menziesii* 211–212
 - Thallopaga* host 250 App.
- Psiguria*, Cucurbitaceae: *Heliconius* butterfly food plants 162–163 & Tab. 1, 165, 167–168 & Fig. 2
- P. spp.* 161–162
 - P. warszewiczii* 162, 167
- Psilaspilates*: lithinine moths on Blechnaceae 250 App.
- Psychoides* (Lepidoptera): fern associations 241 Tab. 1
- psyllids (Aphalaridae-Trioizidae): phytophagous insects 111
- Pteridaceae: lithinine moth hosts 245–246 & Fig. 2, 250 App.
- Pteridium*
- P. aquilinum*: insect food-plants 117
 - P. spp.*: moth associations 241, 245–247 & Fig. 2, 250 App.
- *pteridophagy: lithinine moths on ferns 239–248 & Tabs 1 & Figs 1–2
- arthropod survey 240
- Pteridophyta (ferns): lithinine moths 230–250 & Figs 1–2 & Tabs 1–2 & *see also*
- Pteris* (Pteridaceae): lithinine moth hosts 245–246 & Fig. 2, 250 App.
- "common brake": (*Pteridium*): *Philedia* host association 250 App.
- pupae
- Bicyclus anynana* butterflies: insect prey experiments 17, 19–20 & Fig. 1, 26
 - Pararge aegeria* (speckled wood butterflies) 143, 145–154 & Figs 2–7
 - 'pupal mating': *Heliconius* butterflies 166–167
- purine nucleoside phosphorylase (*Np*, E.C. 2.4.2.1) 72
- Apodemus* 87 App. 2
 - Microtus* 89 App. 2
 - Niviventer* 88 App. 2
- Pyralidae
- (family): fern associations 241 Tab. 1
 - Pyraloidea (superfamily) 241 Tab. 1
 - species-rich Lepidoptera 226
- *quail, Japanese (*Coturnix coturnix japonicus*): genetics of predation 17–21 & Fig. 1
- quokkas (*Setonix brachyurus*): resolution of kangaroo phylogeny using DNA hybridization 309–310, 314 Tab. 1, 316 Tab. 2
- ranch populations: American mink 293–305 & Tabs 1–8 & Figs 1–4
- randomness
- mating: evolutionary genetics of Asian mammals 77
 - tests: isopod *Idotea baltica* experiments 49–50, 55, 58 Tab. 4, 65
- Ranunculaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 119, 126–127 Apps 1–2
- Ranales 119
- rat kangaroos (Potoroinae): resolution of kangaroo phylogeny using DNA hybridization 310
- rufous (*Aepyrymnus rufescens*) 309, 313
- rats *see Niviventer*
- Raunkiaer's life forms 109–111, 121
- *reaction norms: plasticity in *Pararge aegeria* 143–146, 148–150 Figs 2–5
- recruitment rate: novel insect prey into the diet of a bird 17–21 & Fig. 1, 26
- 'red' mimicry rings: *Heliconius* butterflies 162, 164–165 & Tab. 1, 169–170 & Tab. 2, 174–176
- see also H. erato* and *H. melpomene*
- Reecer Creek, Wenatchee Mountains: ants and *Euphilotes enoptes* 211–220 & Figs 2–4 & Tabs 2–5
- regressions
- colour patterns in a lizard 276–278, 282–283 & Tab. 2
 - Euphilotes enoptes* larvae and ants: linear 213, 216
 - fit of flowers around the bodies of bees 373, 375–378 & Tabs 7–8
 - offspring-parent: insect prey experiments 23

- patterns in brood parasitism: linear 263–264 & Fig. 1
- Rensselaerville, New York: fit of flowers around the bodies of bees 361–362 Tabs 1–2, 368, 370 Fig. 7, 373–376 Tabs 5–8
- *reproduction: colour patterns in a lizard 273–276, 280–282 & Tab. 1, 286, 288–290
- reproductive success: ornaments or offspring 251–252, 255–256 & Fig. 2
- potential (PRS) 255–258 & Fig. 2 & Tab. 1
- realized (RRS) 255
- reptiles: sex-linked colour polymorphism 274
- Resedaceae: insect food-plants 112, 114 Fig. 1, 126–127 Apps 1–2
- resistance genes (A2–B2): *Culex pipiens* population structure and genetic invasion 329–330, 340–341
- resource competition: cranial variation in mink 293, 304
- Restoration of Endangered Species, Conceptual Issues, Planning and Implementation*, editors M. L. Bowles and R. J. Whelan (review: Norman W. Moore) 181
- results
- case study of variation in predator preference 104–105 & Tab. 1
 - colour patterns in a lizard 278–285 & Figs 1–6 & Tabs 1–2
 - convergence in harvester ant assemblages 17, 35–39 & Tabs 2–3 & Figs 2–4, 41
 - cranial variation in mink 296–303 & Tabs 3–9 & Figs 1–4
 - Culex pipiens* population structure and genetic invasion 335–339 & Tabs 2–5 & Fig. 2
 - evolutionary genetics of Asian mammals 73–80 & Tabs 1–4 & Figs 2–3
 - genetics of predation 17, 22–24 & Tabs 1–3
 - insect food-plants 109, 112–121 & Tab. 1 & Figs 1–8
 - Madagascan *Tropidophora* 188–200 & Figs 2–8 & Tab. 2
 - ornament growth, variation and dimorphism in *Vidua* 129, 132–138 & Tabs 1–7
 - patterns in species description 227–234 & Figs 1–8 & Tabs 1–4
 - plasticity in *Pararge aegeria* 143, 146–154 & Figs 1–7
 - polymorphism of the isopod *Idotea baltica* 46, 52–61 & Figs 1–6 & Tabs 2–5
 - relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 3–8 & Figs 1–5
 - resolution of kangaroo phylogeny using DNA hybridization 313–321 & Tabs 1–4 & Figs 1–3
 - sexual dimorphism in *Metellina segmentata* 348–350 & Tab. 1 & Figs 2–3
 - unpredictable variation in a facultative mutualism 213–220 & Tabs 1–5 & Figs 1–4
- retentive index: Nelson consensus tree (Madagascan *Tropidophora*) 189
- Rhamnaceae: insect food-plants 112 & Tab. 1, 114 Fig. 1, 126–127 Apps 1–2
- rhinoceros phylogeny: comparative serology 325
- Rhoplocomus* fossil: spicules in Demospongiae 11
- Ribes* (Saxifragaceae): *Thallopaga* associations 247, 250 App.
- rodents
- evolutionary genetics of Asian mammals 69, 71, 73 Tab. 1, 75–77 & Tabs 2–4, 80–81 & Fig. 3, 87–89 App. 2
 - see also *Apodemus*, *Microtus*, *Niniventer*, *Peromyscus* and *Thomomys*
 - laboratory: analogous to ranch mink 305
 - resource limitations 41
- Roger's genetic distances; evolutionary genetics of Asian mammals 72, 75–76 & Tab. 2, 78–79 & Fig. 2
- Rosa*: pollination of trimmed flowers 380
- Rosaceae: insect food-plants 111–112 & Tab. 1, 114 & Fig. 1, 117, 126–127 Apps 1–2
- Rosales 112, 119
- Rosidae 119 & Fig. 5
- Rotuloplocamia*: spicules in Demospongiae 11
- Rubiaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2
- Rubus*: pollination of trimmed flowers 380
- Rumoba* (Dryopteridaceae): *Idiodes* association 250 App.
- Rymättylä: isopod *Idotea baltica* 49–50, 52–57 & Figs 1–3 & Tabs 1–2, 62
- Salicaceae
- insect food-plants 112 Tab. 1, 114 Fig. 1, 126–127 Apps 1–2
 - larval hosts for *Thallopaga* 250 App.
 - Salix* 247, 250 App.
- sampling procedures
- harvester ant data 31–34 & Fig. 1, 36
 - polymorphism of the isopod *Idotea baltica* 46, 49–50, 52–53, 62
- SAN (Arborea, Sardinia): *Culex pipiens* sample site 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2
- sandpipers, spotted: sexual selection acting on females 256 Tab. 1,
- 'saphire': ranch mink pelts 297–299 & Figs 1–2 & Tabs 3–4, 304
- Sapindales: insect food-plants 113
- Saproscincus*: Australian scinclid lizards
- S. challengerii* 274
 - S. galli* 274
 - S. mustelinus* 274
 - S. rosei* 274
- Sardinia: *Culex pipiens* population structure and genetic invasion 329–331, 337–341 & Tabs 3–5
- see also BAL, CFR, CAG1, CAG2, CHI, GER, OLB, SAN, SPR, TEU and VMA
- Sarisa* (lithinine moths): pteridophagy 245–246 & Fig. 2, 250 App.
- SAS: colour pattern polymorphism in a lizard 278
- Saturniidae: pteridophyte associations 240
- *Satyriinae: *Pararge aegeria* (speckled wood butterflies) 143
- sawflies: phytophagous insects 115
- Saxifragaceae
- insect food-plants 114 Fig. 1, 126–127 Apps 1–2
 - larval hosts for *Thallopaga* 250 App.
 - see also *Ribes*
- *scales: unpredictable variation in a facultative mutualism 209–211, 214–216
- Schiziales: lithinine moth hosts 245–246 & Fig. 2, 250 App.
- scinclid lizards: colour patterns 273–275
- see also *Lampropholis delicata* and *Saproscincus*
- Scolytidae (Coleoptera): phytophagous insects 114

- Scotland, Heriot Farm: American mink 299 Fig. 2 & Tab. 4
- screening time: genetics of predation 18, 26
- Scrophulariaceae: insect food-plants 112–114 & Tab. 1 & Fig. 1, 117, 126 127 Apps 1 2
- Scrophulariales 119
- seahorses: sexual selection acting on females 256 Tab. 1,
- seasonality: patterns in brood parasitism 261–262
- *seasonality: plasticity in *Pararge aegeria* 143–146
- seeds
- bait: harvester ant assemblages 29, 31, 33–34, 39–41, 44 App.
 - millet seed 33
 - insect food-plants 111, 122–123
- *selection: colour patterns in a lizard 273
- selection
- fit of flowers around the bodies of bees 355–358, 372–373, 375–380 & Tabs 7 8
 - frequency-dependent: variation in predator preference 93–107 & Figs 1–6 & Tab. 1
 - ornament growth, variation and dimorphism in *Vidua* 129–131, 138, 140
 - polymorphism of the isopod *Idotea baltica* 45 47, 64–66
 - sexual: ornaments or offspring 251 258 & Figs 1–2 & Tab. 1
- *selection lines: genetics of predation 17–20, 24–26 & Fig. 2
- Serbia: *Apodemus agrarius* 72, 78, 85 App. 1
- Sestra* (lithinine moths): pteridophagy 245–246 & Fig. 2, 250 App.
- Selonix*: resolution of kangaroo phylogeny using DNA hybridization 311, 324, 326
- S. brachyurus* (quokkas) 309–310, 314 Tab. 1, 316 Tab. 2, 319 321 & Fig. 2
- *sex-role reversal: sexual selection acting on females 251–252, 254, 257
- *sexual behaviour: butterfly mimicry 160–161, 166–167, 176
- *sexual differences: polymorphism of the isopod *Idotea baltica* 45–68 & Figs 1–6 & Tabs 1–5
- sex-determination 48–49, 62
- see also females and males
- *sexual dimorphism: cranial variation in mink 293–295, 298–301 & Tabs 4–7 & Figs 3–4, 304–305
- sexual dimorphism: *Metellina segmentata* 345–354 & Figs 1–3 & Tab. 1
- *sexual dimorphism: whydahs (*Vidua*) 129–133 & Tab. 1
- *sexual selection: whydahs (*Vidua*) 129–131, 138, 140
- Sexual Selection*, by M. Andersson (review: P. N. Ferns) 92
- Shannon diversity index (calculation of alpha diversity): harvester ant assemblages 34
- *shape: sexual dimorphism in *Metellina segmentata* (PC1) 345–352 & Figs 1–3 & Tab. 1
- Shapiro-Wilk statistic: colour pattern polymorphism in a lizard 285
- *shells: Madagascan *Tropidophora* 183–184, 186, 188–195 & Figs 3–6 & Tab. 2, 198–201–202, 205–208 App
- shorebirds: sexual selection acting on females 256 Tab. 1,
- shrubs: insect food-plants 109, 111, 113, 115, 117–119
- 'silverblu' (SB): ranch mink pelts 297–298–299 & Figs 1–2 & Tabs 3 4, 304–305
- similarity: convergence in harvester ant assemblages 30
- Siproeta stelenis* (nymphalid butterflies): feeding on *Blechnum* (Acanthaceae) 240
- Sirena station, Costa Rica: butterfly behaviour and mimicry in *Heliconius* 164–165 & Fig. 1
- Sitodiplosis mosellana* (wheat blossom midges): staggered diapause 122–123
- *size: sexual dimorphism in *Metellina segmentata* (PC2) 345 352 & Figs 1–3 & Tab. 1
- size
- harvester ant assemblages 29, 34, 36–38 & Tabs 2 3 & Fig. 3
 - predation of isopod *Idotea baltica* 56–58 & Tab. 5, 64–65
 - whydah (*Vidua*) mass 129, 131–138 & Tabs 1–3 & 6–7
- Skjabinylus nasicola* (nematodes) infection: American mink 303
- skulls: variations in mink 293–295 & Tab. 1, 303–304
- skylarks (*Alauda arvensis*) males: flight display compensations 138
- Slatkin's methods
- log scales: *Culex pipiens* migrant numbers 339 & Fig. 2
 - private alleles
 - Culex pipiens* migrant numbers 335, 338–339 & Tab. 5
 - evolutionary genetics of Asian mammals 77 & Tab. 4, 82
- *snails: Madagascan *Tropidophora* 183–208 & Figs 1–8 & Tabs 1 2 & App
- see also acavid, camaenid, polygyrid, *Cerion*, *Ceritium* and *Littorina*
- snakes: colour patterns in a lizard 275, 287
- snout-vent length (SVL): *Lampropholis delicata* 275–278, 280–284 & Tabs 1–2 & Figs 3 & 5
- Society archipelago: dispersal of resistance genes in *Culex pipiens* 339
- see also Moorea and Tahiti 339
- Sod*: Madagascan *Tropidophora* (allozyme data) 186, 189 Fig. 2, 207–208 App.
- Solanaceae
- insect food-plants 112 & Tab. 1, 114 & Fig. 1, 117, 119, 126–127 Apps 1–2
- Solanales 119
- Melinaea* egg-laying hosts 172
- see also *Juanulloa* and *Markea* 172
- Solbrig, O. T., van Ermden, H. M. and van Oordt, P. G. W. J. editors: *Biodiversity and Global Change*. (review: Oliver L. Gilbert) 181–182
- Solenopsis*: harvester ant assemblages
- North America 40
 - South America
 - S. gayi* 44 App.
 - S. granivora* 44 App.
 - S. saevissima* 44 App.
 - S. sp.* 44 App.
- Solothurn, Switzerland: patterns in brood parasitism 263–265 & Fig. 2, 272 App.
- sorbitol dehydrogenase (*Sordh*, E.C. 1.1.1.14.) 72
- evolutionary genetics

- Microtus* 89 App. 2
Niviventer 88 App. 2
 Madagascan *Tropidophora* (allozyme data) 186, 189
 Fig. 2, 207–208 App.
- sources
 Japanese quail: Netherlands 19
 sponges 2–3
 South America
 harvester ant assemblages 29–33 & Fig. & Tab. 1,
 35–41 & Tabs 2–4 & Figs 2–3, 44 App.
see also, Andes, Argentina, Chile, Monte Desert
 and Patagonia
 Neotropics: biogeographic region 226
 Spain
 Demospongiae 3
see also Alboran Island
Pararge aegeria (speckled wood butterflies) 143–147 &
 Fig. 1, 153–154 & Fig. 6
 Sparganiaceae: insect food-plants 114 Fig. 1, 126–127
 Apps 1–2
 Sparganiales 120
 *spatial variation: unpredictable variation in a
 facultative mutualism 209, 214–216, 221
 Spearman rank correlations: ornament growth, variation
 and dimorphism in *Vidua* 132, 136 Tab. 6
 speciation: Taiwanese small mammals 69, 71
 species
 differences: fit of flowers around the bodies of bees
 355–359, 362–363, 368–373 & Figs 6–8 &
 Tab. 5, 377–378
 richness: convergence in harvester ant assemblages
 34, 36–39 & Tabs 2–3
 *species richness: patterns 225–237 & Figs 1–8 &
 Tabs 1–4
 specimens, preserved: *Lampropholis delicata* 273–274,
 276–282 & Figs 1–3 & Tabs 1–2, 288–289
 sphaeroclons (fossils): relationships between polyaxonid
 and monaxonid spicules in Demospongiae 1, 3,
 7–12 & Figs 4 & 7
 spicules: relationships between polyaxonid and
 monaxonid in Demospongiae 1–15 & Figs 1–7
see also monaxonid, polyaxonid and tetraxonid
 spiders
 habitat structure 122
 sexual dimorphism in *Metellina segmentata* 345–352 &
 Figs 1–3 & Tab. 1
see also orb-weaving
 spines: relationships between polyaxonid and
 monaxonid spicules in Demospongiae 2–7, 12
 Spirophorida: spicules in Demospongiae 12
 sponges: relationships between polyaxonid and
 monaxonid spicules in Demospongiae 1–15 &
 Figs 1–7
 SPR (San Priamo, Sardinia): *Culex pipiens* sample site
 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2
 standard deviations: fit of flowers around the bodies of
 bees 361–363 & Tab. 3, 377
 standard errors (means): fit of flowers around the
 bodies of bees 373 Tab. 5
 StatGraphics (STSC): ornament growth, variation and
 dimorphism in *Vidua* 132
 Stathmopodinae (lepidopteran subfamily): fern
 associations 241 Tab. 1
 statistical procedures
 colour pattern polymorphism in a lizard 276–285, 287
see also chi-square, G, Shapiro-Wilk, likelihood
 ratio, *t* and Wilcoxon tests 276–280
 convergence in harvester ant assemblages 34–35, 75
see also means and Shannon diversity index
 cranial variation in mink 293, 302
see also Cohens Kappa, computer programs,
 Mahalanobis distances, Mann Whitney,
 measurement errors, principal components
 (PCs) and Wilks' lambda
Culex pipiens population structure and genetic
 invasion 335–336
see also Bonferroni method, correlation coefficient,
 D, *F*, Fisher's tests, Haldane's exact test,
 Hardy-Weinberg, Markov chain and
 Slatkin's methods 333
 evolutionary genetics of Asian mammals *see*
 bootstrap, chi-square, F-statistics, genetic
 distances and Slatkin's methods
 facultative mutualism between *Euphilotes enoptes*
 larvae and ants: *see* Morisita's index, G and
t-tests
 fit of flowers around the bodies of bees 357, 361,
 372, 373–375 Tabs 5–7
see also means, correlations, standard deviations
 and errors, and variances
 genetics of predation 22–23 & Tab. 1
see also correlations, Kruskal-Wallis,
 Mann-Whitney and regressions
 insect food-plants 110–112, 123
 Czekanowski coefficient 111–112
 ornament growth, variation and dimorphism in
Vidua 132
see also Bonferroni adjustments, coefficients of
 variation, covariance, F-ratios,
 Kolmogorov-Smirnov tests, principal
 components, Spearman rank correlations,
 Statgraphics (STSC) 132
 patterns in brood parasitism: correlations,
 Kolmogorov-Smirnov test and Tukey Kramer
 methods 262
 plasticity in *Pararge aegeria* 152
 polymorphism of the isopod *Idotea baltica* 52–57 &
 Tabs 2–3
see also chi-square, heterogeneity (G), randomness,
 sampling, and Student's *t* test
 resolution of kangaroo phylogeny 309, 312–313,
 318–325 & Figs 2–3
 bootstrap and jackknife methods 309, 312–313,
 318–325 & Figs 2–3
see also CONSENSE, FITCH and PHYLIP
 sexual dimorphism in *Metellina segmentata*: *t* tests
 347–350 & Tab. 1 & Fig. 3
see also analyses, computer programs, indices and
 regressions
 Sterrhinae: patterns in species description 227–228 &
 Figs 1–2, 232–233 & Fig. 6 & Tabs 3–4
 Stevens, Peter F. *The Development of Biological Systematics*.
 (review: David Williams) 386
 Sthenurinae: *Lagostrophus fasciatus* classification 310
 sticklebacks: sexual selection acting on females 256
 Tab. 1
 stigmas: fit of flowers around the bodies of bees
 355–358, 360–367 & Figs 2–4, 369–375 & Fig. 6
 & Tab. 5, 378–380
 stigma-anther separation 379

- stoats: quantitative genetics 294
- Strepera graculina* (currawongs): lizard predators 287
- stripes, white mid-lateral: *Lampropholis delicata* 273–289 & Figs 1–4 & Tabs 1–2
- structures, genetic population: small mammals in Taiwan 69
- Student's *t* tests *see t*-tests
- study sites
- convergence in harvester ant assemblages 31–33 & Fig. 1
 - unpredictable variation in a facultative mutualism 211–213
- Stylidium*: pollination success 378
- *subarid ecosystems: patterns in brood parasitism 261
- subtropics: patterns in brood parasitism 261–262
- Sumatra: *Niviventer* rats 81
- summaries: evolutionary genetics of Asian mammals 73–75
- summer: plasticity in *Pararge aegeria* 143–153 & Fig. 2
- sunbirds (*Nectarinia johnstoni*): sexual signalling 139
- superoxide dismutase (*Sod-1*, *Sod-2*), E.C. 1.15.1.1) 72
- Sod-1*
- Apodemus* 87 App. 2
 - Microtus* 89 App. 2
 - Niviventer* 88 App. 2
- survival: colour patterns in a lizard 273–275, 277, 279–280, 283–284 & Fig. 4 & Tab. 2, 286–289
- swallows: sexual selection acting on females 256 Tab. 1
- barn (*Hirundo rustica*)
- male European 258
 - male North American 258
- Sweden: *Pararge aegeria* (speckled wood butterflies) 145, 152–153
- central 143–145, 147 & Fig. 1, 149–152 & Fig. 3
 - southern 143–148 & Figs 1–2, 149–152 & Fig. 4
- Switzerland: patterns in brood parasitism 263–265 & Fig. 2, 272 App.
- see also* Solothurn
- Sydney: colour pattern polymorphism in a lizard 275–277, 279–280 & Figs 1–2
- sympatry: *Impatiens capensis* and *I. pallida* 355, 371–372 & Fig. 8, 377
- *synonyms: patterns in species description 225–226, 229–236 & Figs 3–8 & Tabs 3–4
- synonyms: *Tropidophora* (Madagascan snails) 184, 202
- systematics: Madagascan *Tropidophora* 183, 185–186
- t*-tests
- colour patterns in a lizard 276
 - one-tailed: estimates of Morisita's index 213
 - polymorphism of the isopod *Idotea baltica* 56–58 & Tab. 5, 56–58 & Tab. 5
 - sexual dimorphism in *Metellina segmentata* 347–350 & Tab. 1 & Fig. 3
- Tacparia*: secondary shift from ferns to seed plants 247, 250 App.
- Tahiti, Society archipelago: dispersal of resistance genes in *Culex pipiens* 339
- tails
- African whydahs (*Vidua*) 129–141 & Tabs 1–7
 - Lampropholis delicata* 275–276, 286
 - regenerated 273, 276, 279, 286
 - streamers: *Hirundo rustica* (barn swallows) 258
- Taiwan: evolutionary genetics of Asian mammals 69–89 & Figs 1–3 & Tabs 1–4 & Apps 1–2
- see also* Mt Guarshan and Mt Yushan
- Tamaricaceae: insect food-plants 114–115 & Fig. 1, 117, 126–127 Apps 1–2
- Tapinoma sessile* (Dolichoderinae): tending *Euphilotes enoptes* larvae 213 Tab. 1
- Tarpiscan Creek: Wenatchee Mountains 212–220 & Figs 1–4 & Tabs 2–5
- tarsi: African whydahs (*Vidua*) 131, 133–134 & Tabs 1–3, 137–138 Tabs 6–7
- Tasmania (T): *Lampropholis delicata* 275–276, 279 Fig. 1
- Taxaceae: insect food-plants 114–115 & Fig. 1, 126–127 Apps 1–2
- taxonomy: insect food-plants 109–113, 117–123 & Figs 3–8
- Tc-6*: Madagascan *Tropidophora* (allozyme data) 186
- Teb-9*: Madagascan *Tropidophora* (allozyme data) 186
- Teichobiinae (lepidopteran subfamily): fern associations 241 Tab. 1
- *temperature: colour patterns in a lizard 273, 285–286 & Fig. 6, 100
- temperatures: plasticity in *Pararge aegeria* 144, 146, 153
- *temporal variation: unpredictable variation in a facultative mutualism 209, 214–216, 221
- Tenebrio molitor* (light brown mealworms): insect prey experiments 19
- Terpioz*: relationships between polyaxonid and monaxonid spicules in Demospongiae 11
- T. hoshimoto* 3
- Tertiary sponge-beds: relationships between polyaxonid and monaxonid spicules in Demospongiae 3, 7–8 & Fig. 4
- Tethyan relict: relationships between polyaxonid and monaxonid spicules in Demospongiae 12
- Tetractinomorpha (subclass): Demospongiae 12
- Tetramorium*: Australian harvester ant assemblages 40
- tetragonid spicules: relationships between polyaxonid and monaxonid in Demospongiae 1–2, 12
- TEU (Teleuda, Sardinia): *Culex pipiens* sample site 332–334 & Tab. 1 & Fig. 1, 336 Tab. 2
- Thallophaga*: secondary shift from ferns to seed plants 247, 250 App.
- The Development of Biological Systematics*, by Peter F. Stevens. (review: David Williams) 386
- Thelypteridaceae: *Ischalis* host association 244–245 Tab. 2 & Fig. 2, 250 App.
- Thelypteris* (*Petrophora*) 250 App.
- see also* *Mesophlebia* and *Pneumatopteris*
- Theodoxus fluviatilis*: epibiont background for isopod *Idotea baltica* 48
- therophytes (seeds: annuals): insect food-plants 111–112, 115–116 & Fig. 2
- thiarid *Pachymelania*: shell variation 202
- Thomomys bottae* (pocket gophers)
- gene flow 82
 - genetic structure 70
- Thuja*: *Thallophaga* host 250 App.
- Thylogale* (pademelons): resolution of kangaroo phylogeny using DNA hybridization 309–311, 315, 318–319, 324, 326
- T. stigmatica* 313–314 Tab. 1, 317 Tab. 2, 320–321 Fig. 2
- T. thetis* 313–316 Tabs 1–2 & Fig. 1, 320–321 Fig. 2
- Thylogale-Petrogale-Dendrolagus* clade 319
- Thymelaeaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2

- 'tiger' mimicry rings: butterflies 162, 164–165 & Fig. 1, 169–170 & Tab. 2, 172, 175–177
 see also *Heliconius hecale*, *H. ismenius*, *Melinaea scylax* and *Tithorea tarricina*
- Tiliaceae: insect food-plants 114 Fig. 1, 119, 126–127 Apps 1–2
- Timea: spicules in Demospongiae 12
- Times Atlas of the World (vegetative classes): patterns in brood parasitism 262–264 & Tab. 1, 266 Fig. 3, 270–272 App.
- timing
 genetics of predation
 FAST (shortest) and SLOW (longest PRT) 19–20, 22–27 & Tab. 1
 prey recruitment (PRT) 18–24 & Tab. 2
 see also selection lines
 ornament growth, variation and dimorphism in *Vidua* 129–132, 135–136 & Tab. 4, 139–140
- Tineidae (lepidopteran family): fern associations 241 Tab. 1
- Tineoidea (lepidopteran superfamily): fern associations 241 Tab. 1
- Tithorea tarricina*: butterfly behaviour and mimicry 164–165 & Fig. 1, 172
- Tortricidae: pteridophyte associations 240
- Townsville, North: *Lampropholis delicata* 276, 279–280 Figs 1–2
- toxicities: genetics of predation 17–19, 21, 23, 26
 see also seven-spot ladybirds (*Coccinella septempunctata*)
- traits: sexual selection acting on females 251–259 & Figs 1–2 & Tab. 1
- transects (1–3): Taiwan 71–73, 76–78 & Tab. 4 n., 82, 85–86 App. 1
- treatment sequence: insect prey experiments 20–21 & Fig. 1
- tree kangaroos (*Dendrolagus*): resolution of kangaroo phylogeny using DNA hybridization 310–311, 319, 325
D. matschieri 309
- *tree snails: Madagascan *Tropidophora* 183
- trees
 consensus
 50% majority rule: evolutionary genetics of Asian mammals 78, 80 Fig. 3
 Nelson: Madagascan *Tropidophora* 189
 phylogenetic: resolution of kangaroo phylogeny using DNA hybridization 312–313, 318, 323–326
- trees (plants)
 gum (*Eucalyptus* spp.): *Lampropholis delicata* experimental areas 277
 insect food-plants 109, 111, 113–115, 117, 119, 122
Trichosturus: resolution of kangaroo phylogeny using DNA hybridization 309, 313, 316–317 Tab. 2, 319–321 & Fig. 2, 323
T. vulpecula (brush-tailed phalanger) 309, 313–314 & Tab. 1, 316–317 Tab. 2, 320–321 Fig. 2
- Trikettrion*: spicules in Demospongiae 11–12
- Triozidae (psyllids): phytophagous insects 111
- tropics: patterns in brood parasitism 261–262, 266
- Tropidophora*: Madagascan snails 183–208 & Figs 1–8 & Tabs 1–2 & App
T. balteata (synonym: *T. unicarinata*) 184
T. bemaraensis (synonym: *T. tenuis*) 184
T. besalampiensis (synonym: *T. tenuis*) 184
T. bicarinata (ANSP identification) 201
T. castanea (synonym: *T. tenuis*) 184
T. cavernarum (synonym: *T. tenuis*) 184
T. chavani (ANSP) 201
T. crenulata (synonym: *T. tenuis*) 184
T. deburghie (synonym: *T. unicarinata*) 184
T. denisi (synonym: *T. tenuis*) 184
T. formosa (synonym: *T. unicarinata*) 184
T. johnsoni (synonym: *T. tenuis*) 184
T. lineata (synonym: *T. tenuis*) 184
T. microchasma (synonym: *T. tenuis*) 184
T. oppessulata (synonym: *T. tenuis*) 184
T. perfecta (synonym: *T. unicarinata*) 184
T. principalis (ANSP identification) 201
T. reesi (synonym: *T. tenuis*) 184
T. sarodranensis (synonym: *T. tenuis*) 184
T. semidecussata (ANSP) 201
T. soulaiana (synonym: *T. unicarinata*) 184
T. tenuis (ANSP) 184, 201
T. tomlini (synonym: *T. tenuis*) 184
T. tricarinata (ANSP) 184, 201
T. t. bicarinata 184
T. t. tricarinata 184
T. tulearensis (synonym: *T. tenuis*) 184
T. unicarinata (ANSP) 184, 201
T. vignalii (synonym: *T. tenuis*) 184
- Trout Creek, New York: fit of flowers around the bodies of bees 361–362 Tabs 1–2, 368–369 & Fig. 6, 373–376 Tabs 5–8
- Tsuga*: *Thallophaga* host 250 App.
- tubercles: relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 3, 5 & Fig. 2, 9 Fig. 6
- Tukey Kramer method: patterns in brood parasitism 262
- Tukey multiple comparison: pollen removal and deposition on *Impatiens capensis* 367 Fig. 5
- turkeys, brush-: sexual selection acting on females 256 Tab. 1,
- tylostyles, tuberos: relationships between polyaxonid and monaxonid spicules in Demospongiae 1, 3, 5 Fig. 2, 9 Fig. 4, 11–12
- Typhaceae: insect food-plants 114 Fig. 1, 126–127 Apps 1–2
- Ulmaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 119, 126–127 Apps 1–2
- Umbelliferae: insect food-plants 112 Tab. 1, 114 Fig. 1, 126–127 Apps 1–2
- United States (USA)
 ants and *Euphilotes enoptes* larvae 209, 211
 harvester ant assemblages 32 Fig. 1
 land snails 200
 see also *Mesodon*
Microtus californicus 72, 86 App. 1
 patterns in brood parasitism 262, 270 App.
 see also New York
- univariate analyses: sexual dimorphism in *Metellina segmentata* 345–347, 350–351
 see also *t*-tests
- UPGMA clustering: evolutionary genetics of Asian mammals 72, 77–79 & Fig. 2, 81–82
- Urosaurus ornatus*: sex-linked colour polymorphism 274
- Urticaceae: insect food-plants 112 Tab. 1, 114 Fig. 1, 119, 126–127 Apps 1–2
- Urticales 118–119

- Uta stansburiana*: sex-linked colour polymorphism 274
- Valerianaceae: insect food-plants 114 Fig. 1, 126–127
Apps 1–2
- valid names: patterns in species description 225–226,
229, 231–235 & Tab. 4 & Fig. 8
- variances
fit of flowers around the bodies of bees 359,
363–364 & Tab. 4, 375 Tab. 7, 378
see also ANOVAs and MANOVAs
- *variation: predator preferences 93–107 & Figs 1–6 &
Tab. 1
- variations
colour patterns in a lizard 273–275, 277–279 & Fig.
1, 281, 285
convergence in harvester ant assemblages 30, 33,
39–40
coefficient 39–40
cranial in mink 293–306 & Tabs 1–9 & Figs 1–4
evolutionary genetics of Asian mammals
allozymic 73–75
genetic 70, 74–75, 81–83
plasticity in *Pararge aegeria* 143–144, 151
polymorphisms of isopod *Idotea baltica* 45–48, 52–56,
61–63, 65–66
rate of recruitment of novel insect prey into quail
diet 17–18
sexual dimorphism in *Metellina segmentata* 345,
348–351
unpredictable variation in a facultative mutualism
209–211, 214–221
see also spatial and temporal
- vegetative classes: patterns in brood parasitism
262–267 & Tab. 1
- Verbenaceae: insect food-plants 114 Fig. 1, 126–127
Apps 1–2
- Vetulina* (sponge fossils): New Zealand 1, 3, 7 Fig. 4, 10
V. oamaruensis 3
- Viburnum*: insect food-plants
V. lentana 116
V. opulus 116
V. tinus 116
- **Vidua paradisaea*: ornament growth, variation and
dimorphism 129–141 & Tabs 1–7
- **Vidua regia*: ornament growth, variation and
dimorphism 129–141 & Tabs 1–7
Vidua (whydahs) 129, 139
- Viola cazortensis*: pollination by hawkmoths 379
- Violaceae: insect food-plants 114 Fig. 1, 126–127
Apps 1–2
- Violales 119
- Vitaceae: insect food-plants 112 Tab. 1, 114–115 &
Fig. 1, 117, 126–127 Apps 1–2
- VMA (Villamalla, Sardinia): *Culex pipiens* sample site
332–334 & Tab. 1 & Fig. 1, 336–338 & Tab. 2
- voles: evolutionary genetics of Asian mammals see
Microtus kikuchii
- Wallabia*: resolution of kangaroo phylogeny using
DNA hybridization 309, 311, 319, 325–326
W. bicolor (swamp wallabies) 309–310, 312–314 &
Tab. 1, 317–318 Tabs 2–4, 320–321 Fig. 2, 325
- wallabies: resolution of kangaroo phylogeny using
DNA hybridization 309, 324
- banded hare- (*Lagostrophus fasciatus*) 310, 326
- forest (*Dorcopsulus vanheurni*) 309, 311, 313, 323
- hare (*Lagorchestes*) 310, 326
- nail-tailed (*Onychogalea*): classification 310, 326
- Parry's (*Macropus parryi*) 309, 312, 314 Tab. 1,
317–321 & Tabs 2–3, 325
- red-necked (*Macropus rufogriseus*) 309, 314–315 Tab.
1 & Fig. 1, 317–319 & Tabs 2–4, 325
- rock (*Petrogale*) 309–311
- swamp (*Wallabia bicolor*) 309–310, 312
- wallaroos: resolution of kangaroo phylogeny using
DNA hybridization: *Macropus robustus* 309,
318–319 & Tabs 3–4
- *warning colouration: butterfly behaviour and
mimicry 159–161
- *warning colouration: genetics of predation 17–18
- Washington state: ants and *Euphilotes enoptes* larvae
209, 211
- waterboatmen, female (*Notonecta*): variation in predator
preference 93, 104–106 & Fig. 6
- weasels: quantitative genetics 294
- web builders: sexual dimorphism in *Metellina segmentata*
345–346, 352
- weeds: insect food-plants 109, 112, 117
- weevils: phytophagous insects 122
- Weislo, WT: patterns in brood parasitism 262–263, 266
- Weld Preserve: Long Island: fit of flowers around the
bodies of bees 361–362 & Tabs 1–2, 373–376
Tabs 5–8
- Wenatchee Mountains, Washington state: ants and
Euphilotes enoptes 211–212
see also Reecer and Tarpiscan Creeks
- Western Palaearctic: biogeographic region (3) 226,
228–232 & Tabs 1–2 & Fig. 3, 234–235 & Fig. 8
see also Europe
- whydahs (*Vidua*): ornament growth, variation and
dimorphism 129–141 & Tabs 1–7
paradise see *V. paradisaea*
shaft-tailed see *V. regia*
- widows
Euplectes
male flight display compensations 138
ornament length: *E. prognus* 139
sexual selection acting on females 256 Tab. 1
- Wilcoxon 2-sample test: colour pattern polymorphism
in a lizard 284–285
- Wilk's lambda
colour pattern polymorphism in a lizard 281 &
Tab. 1
cranial variation in mink 297–298, 302
- wings: African whydahs (*Vidua*) 131, 133–135 & Tabs
1–3, 136–138 & Tabs 6–7
- winter: plasticity in *Pararge aegeria* 143–148 & Fig. 2,
150, 152–153
- wintergreen species: insect food-plants 116
- woody plants: insect food-plants 109, 111, 113–117,
121–123
see also shrubs and trees
- Xanthorrea* spp. (grass trees): *Lampropholis delicata*
experimental areas 277
- 'yellow' mimicry rings: *Heliconius* butterflies 164–165 &
Fig. 1, 167–170 & Fig. 2 & Tab. 2, 175}